

# Shadowing and the Viability Kernel Algorithm

Janosch Rieger\*

May 30, 2008

## Abstract

The aim of this paper is to derive estimates for the accuracy of the viability algorithm for systems which have shadowing properties. Recently developed shadowing results are applied in order to prove that the algorithm is linearly convergent for a certain class of right hand sides.

**Key words.** viability kernel algorithm, shadowing  
**AMS(MOS) subject classifications.** 34A60, 37C50

## 1 Introduction

Viability kernels of differential inclusions are of considerable interest, because many theoretical and practical problems can be reformulated as viability problems. A standard reference on viability theory is the book [2] by Aubin. In [5], Frankowska and Quincampoix proposed a first algorithm for the computation of viability kernels, and Saint-Pierre succeeded to prove the convergence of a fully discretized and hence implementable algorithm in [11]. This viability kernel algorithm was later generalized to impulsive differential inclusions in [4].

The shadowing property has been thoroughly investigated in the framework of dynamical systems, see [7] and [8]. In set-valued systems, shadowing theory is still under development. First attempts to generalize the classical ideas can be found in [1], [6], [9], [10], and [12].

As viability theory describes the behaviour of trajectories on the unbounded time interval  $[0, \infty)$ , it seems natural to use shadowing results as tools in this context. The aim of this paper is to derive estimates for the

---

\*Supported by the Hausdorff Research Institute for Mathematics, Bonn.

accuracy of the fully discretized viability algorithm for systems which have the shadowing property. In Section 4, a shadowing theorem from [10] is applied in order to show that the viability algorithm is linearly convergent for a certain class of differential inclusions.

## 2 Definitions

Let  $X$  be a finite-dimensional vector space with unit ball  $B \subset X$ , and let  $K \subset X$  be a compact subset. By  $\text{dist}(\cdot, \cdot)$  and  $\text{dist}_H(\cdot, \cdot)$  we will denote the one-sided and the symmetric Hausdorff distance, respectively.

Consider the autonomous differential inclusion

$$\dot{x}(t) \in F(x(t)) \text{ for almost every } t \geq 0, \quad x(0) = x_0 \in X \quad (1)$$

and its  $\rho$ -flow

$$G_\rho : X \rightrightarrows X, \quad x \mapsto \mathcal{R}(0, \rho, x), \quad (2)$$

where  $\mathcal{R}(0, \rho, x)$  denotes the reachable set of (1) at time  $\rho$ . For any  $h > 0$  let  $X_h \subset X$  be a countable subset with

$$\forall x \in X, \exists x_h \in X_h \text{ with } |x - x_h| \leq \alpha(h) \quad (3)$$

and  $\lim_{h \rightarrow \infty} \alpha(h) = 0$ . Given any subset  $A \subset X$  and  $\varepsilon > 0$ , we define

$$A^\varepsilon := A + \varepsilon B \text{ and } A_h^\varepsilon := A^\varepsilon \cap X_h. \quad (4)$$

Consider the semi-discretized Euler scheme

$$\Gamma_\rho : X \rightrightarrows X, \quad x \mapsto x + \rho F(x) \quad (5)$$

and the fully discretized scheme

$$\Gamma_{\rho, h} : X_h \rightrightarrows X_h, \quad x_h \mapsto (x_h + \rho F(x_h) + k(\rho, h)B) \cap X_h, \quad (6)$$

with  $k(\rho, h) := (2 + L\rho)\alpha(h)$ , where  $L > 0$  will be a Lipschitz constant of a restriction of  $F$  (see assumption (iii) below).

**Definition 1.** Let  $G : X \rightrightarrows X$  be a set-valued mapping. Every sequence  $(p_n)_{n \in \mathbb{N}}$  with  $p_{n+1} \in G(p_n)$  for all  $n \in \mathbb{N}$  is called an orbit of  $G$ . A sequence  $(x_n)_{n \in \mathbb{N}}$  satisfying  $\text{dist}(x_{n+1}, G(x_n)) \leq d$  for every  $n \in \mathbb{N}$  is called a  $d$ -pseudotrajectory of  $G$ .

**Definition 2.** A mapping  $G$  has the  $(d, \varepsilon)$ -shadowing property in  $K$ , if for every  $d$ -pseudotrajectory  $(x_n)_{n \in \mathbb{N}} \subset K$  there is an orbit  $(p_n)_{n \in \mathbb{N}}$  of  $G$  such that  $|x_n - p_n| \leq \varepsilon$  for all  $n \in \mathbb{N}$ .

**Definition 3.** A mapping  $G$  has the inverse  $(d, \varepsilon)$ -shadowing property in  $K$ , if for every continuous mapping  $\Phi : X \rightrightarrows X$  with compact and convex values satisfying

$$\text{dist}_H(\Phi(x), G(x)) \leq d$$

for all  $x$  in a neighbourhood of  $K$  and every orbit  $(p_n)_{n \in \mathbb{N}} \in K$  of  $G$  there is an orbit  $(x_n)_{n \in \mathbb{N}}$  of  $\Phi$  such that  $|p_n - x_n| \leq \varepsilon$  for all  $n \in \mathbb{N}$ .

The definition of the inverse shadowing property is kept slightly vague on purpose, because the required size of the neighbourhood depends on the shadowing theorem which is applied in order to ensure that the system has this property (see Section 4 for a concrete example).

**Definition 4.** A subset  $D \subset X$  is a discrete viability domain of a set-valued mapping  $G$ , if

$$G(x_0) \cap D \neq \emptyset \text{ for all } x_0 \in D.$$

The discrete viability kernel  $\text{Viab}_G(K)$  is the largest closed discrete viability domain contained in  $K$ .

Analogously, a subset  $D \subset X$  is a viability domain of the differential inclusion (1), if for any  $x_0 \in D$  there exists a solution  $x : [0, \infty) \rightarrow D$  such that  $x(0) = x_0$ . The viability kernel  $\text{Viab}_F(K)$  is the largest closed viability domain of (1) contained in  $K$ .

Under mild assumptions on  $F$  and  $G$ , both types of viability kernels are well-defined, compare e.g. [2].

The viability kernel algorithm presented in [11] computes the viability kernel  $\text{Viab}_{\Gamma_{\rho,h}}(K_h^\varepsilon)$  for a suitable  $\varepsilon > 0$  instead of the analytical object  $\text{Viab}_F(K)$ . As  $K^\varepsilon$  is compact,  $K_h^\varepsilon$  is a finite set, and thus the algorithm terminates after at most  $\#K_h^\varepsilon$  steps. Estimating the accuracy of the algorithm amounts to estimating the Hausdorff distance  $\text{dist}_H(\text{Viab}_F(K), \text{Viab}_{\Gamma_{\rho,h}}(K_h^\varepsilon))$ . Please note that the mappings  $\Gamma_\rho$ ,  $\Gamma_{\rho,h}$ , etc. are defined in a slightly different way in the present paper for technical reasons, but that the basic idea has been adopted without any changes.

Throughout this paper, we will suppose the following assumptions:

- (i) The viability kernel  $\text{Viab}_F(K)$  is stable in the sense that there exists an  $\varepsilon_0 > 0$  such that  $\text{Viab}_F(K) = \text{Viab}_F(K^{\varepsilon_0})$ .
- (ii) There exist a  $d_0^{(s)} > 0$  and a  $d_0^{(is)} > 0$  such that the  $\rho$ -flow  $G_\rho$  has
  - (iia) the  $(d, \varphi(d))$ -shadowing property in  $K^{\varepsilon_0}$  for  $d \in (0, d_0^{(s)}]$  and

(iib) the inverse  $(d, \psi(d))$ -shadowing property in  $K$  for  $d \in (0, d_0^{(is)}]$ ,

where  $\varphi, \psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  are increasing functions with  $\lim_{d \rightarrow 0} \varphi(d) = 0$  and  $\lim_{d \rightarrow 0} \psi(d) = 0$ .

(iii) The mapping  $F$  is Lipschitz-continuous in  $K^{\varepsilon_0}$  with Lipschitz constant  $L > 0$  and has compact and convex values.

## 3 Results

### 3.1 General estimates

**Observation 1.** *For any compact set  $A \subset X$ , the viability kernels can be characterized by*

$$\text{Viab}_G(A) = \{x_0 \in A : \exists (p_n)_{n \in \mathbb{N}} \subset A \text{ with } p_0 = x_0 \text{ and } p_{n+1} \in G(p_n) \forall n \in \mathbb{N}\}$$

and

$$\text{Viab}_F(A) = \{x_0 \in A : \exists \text{ a solution } x : [0, \infty) \rightarrow A \text{ of (1) with } x(0) = x_0\}.$$

*It is obvious that the right hand sides are the largest viability domains contained in  $A$ . Under mild assumptions on  $F$  and  $G$  they are closed, compare e.g. Theorem 3.5.3 in [2].*

**Observation 2.** *Because of assumption (iii),  $F$  is bounded on  $K^{\varepsilon_0}$  by  $\|F\|_\infty = M < \infty$ . Thus any solution  $x$  of (1) remaining in  $K^{\varepsilon_0}$  satisfies*

$$|x(t) - x(0)| \leq \int_0^t |\dot{x}(s)| ds \leq Mt. \quad (7)$$

*If  $x(0) \in K^\varepsilon$  with  $0 < \varepsilon < \varepsilon_0$ , we have  $x(t) \in K^{\varepsilon_0}$  for all  $t \in [0, \rho]$  with  $0 < \rho < \frac{\varepsilon_0 - \varepsilon}{M}$ . Otherwise  $0 < t_0 := \inf\{t \in [0, \rho] : x(t) \notin K^{\varepsilon_0}\} < \rho$  and (7) holds for all  $0 \leq t \leq t_0$ . But then*

$$|x(t_0) - x(0)| \leq Mt_0 < M\rho \leq \varepsilon_0 - \varepsilon$$

*implies that  $x(t_0)$  is in the interior of  $K^{\varepsilon_0}$ , which is a contradiction. Thus we can apply (7) for all  $t \in [0, \rho]$ . If  $x(0) \in K^\varepsilon$  and  $x(\rho) \in K^\varepsilon$ , we can apply it forwards in time from  $x(0)$  and backwards in time from  $x(\rho)$  in order to obtain*

$$\text{dist}(x(s), K^\varepsilon) \leq \frac{1}{2}M\rho \quad \forall s \in [0, \rho]. \quad (8)$$

**Lemma 3.** *We have*

$$\text{Viab}_F(K) = \text{Viab}_{G_\rho}(K^\varepsilon)$$

for all  $0 \leq \varepsilon < \varepsilon_0$  and  $0 < M\rho < \varepsilon_0 - \varepsilon$ .

*Proof.* Obviously  $\text{Viab}_F(K) \subset \text{Viab}_{G_\rho}(K^\varepsilon)$ . But

$$\begin{aligned} x_0 \in \text{Viab}_{G_\rho}(K^\varepsilon) &\Rightarrow \exists (p_n)_{n \in \mathbb{N}} \subset K^\varepsilon : p_0 = x_0, p_{n+1} \in G_\rho(p_n) \forall n \in \mathbb{N} \\ &\Rightarrow \exists \text{ a solution } x : [0, \infty) \rightarrow X \text{ of (1) : } x(n\rho) = p_n \in K^\varepsilon \\ &\Rightarrow \text{dist}(x(t), K^\varepsilon) \leq \frac{1}{2}M\rho \forall t \geq 0 \\ &\Rightarrow x_0 \in \text{Viab}_F(K^{\varepsilon + \frac{1}{2}M\rho}) = \text{Viab}_F(K) \end{aligned}$$

by assumption (i) and Observation 2. □

The following Lemma is contained implicitly in many works, because it estimates the local error of the semi-discretized Euler-scheme. Nevertheless it is included explicitly in this paper for the sake of self-containedness and readability.

**Lemma 4.** *The error of approximation between  $G_\rho$  and  $\Gamma_\rho$  is*

$$\text{dist}_H(G_\rho(x_0), \Gamma_\rho(x_0)) \leq M\rho(e^{L\rho} - 1)$$

for all  $0 < \varepsilon < \varepsilon_0$ ,  $x_0 \in K^\varepsilon$ , and  $\rho > 0$  such that  $M\rho e^{L\rho} \leq \varepsilon_0 - \varepsilon$ .

*Proof.* Let a solution  $x : [0, \rho] \rightarrow X$  of (1) with  $x(0) = x_0$  be given. Because of (7),  $x(s) \in K^{\varepsilon_0}$  for all  $s \in [0, \rho]$ . As  $F$  has convex values, we can identify the Euler-step  $\Gamma_\rho(x_0)$  with the reachable set of the constant differential inclusion

$$\dot{e}(t) \in F(x_0), \quad e(0) = x_0 \tag{9}$$

at time  $\rho$ . Since

$$\text{dist}(\dot{x}(t), F(x(0))) \leq \text{dist}(F(x(t)), F(x(0))) \leq L|x(t) - x(0)| \leq LMt,$$

the Filippov theorem (cf. Theorem 2.4.1 in [3]) guarantees the existence of a solution  $e : [0, \rho] \rightarrow X$  of (9) satisfying

$$|x(t) - e(t)| \leq \int_0^t e^{L(t-s)} LMt ds = Mt(e^{Lt} - 1)$$

for all  $t \in [0, \rho]$ , and in particular

$$\text{dist}(x(\rho), \Gamma_\rho(x_0)) \leq |x(\rho) - e(\rho)| \leq M\rho(e^{L\rho} - 1).$$

Conversely, let  $\eta \in F(x_0)$  be given and consider the corresponding linear trajectory  $e(t) := x_0 + t\eta$  for  $t \in [0, \rho]$  of  $\Gamma_\rho$ . As

$$\text{dist}(\dot{e}(t), F(e(t))) \leq \text{dist}(F(x_0), F(e(t))) \leq LMt,$$

the Filippov theorem yields a solution  $x : [0, \rho] \rightarrow X$  of (1) with  $x(0) = x_0$  and

$$|x(t) - e(t)| \leq \int_0^t e^{L(t-s)} LMt ds = Mt(e^{Lt} - 1)$$

for all  $t \in [0, \rho]$ , and in particular

$$\text{dist}(e(\rho), G_\rho(x_0)) \leq |e(\rho) - x(\rho)| \leq M\rho(e^{L\rho} - 1).$$

□

### 3.2 Estimates using the shadowing and the inverse shadowing property

**Lemma 5.** *If  $M\rho(e^{L\rho} - 1) \leq d_0^{(is)}$ ,  $\varepsilon_1 := \psi(M\rho(e^{L\rho} - 1)) \leq \varepsilon_0$  and  $M\rho e^{L\rho} \leq \varepsilon_0$  then*

$$\text{dist}(\text{Viab}_F(K), \text{Viab}_{\Gamma_\rho}(K^{\varepsilon_1})) \leq \varepsilon_1,$$

*Proof.* Let  $p_0 \in \text{Viab}_F(K) = \text{Viab}_{G_\rho}(K)$  be given. Hence there exists an orbit  $(p_n)_{n \in \mathbb{N}}$  of  $G_\rho$  such that  $p_n \in K$  for all  $n \in \mathbb{N}$ . As  $\Gamma_\rho$  is continuous with compact and convex values, Lemma 4 ensures that  $\Gamma_\rho$  is an approximation of  $G_\rho$  in the sense of assumption (iib), which in turn yields the existence of an orbit  $(x_n)_{n \in \mathbb{N}}$  of  $\Gamma_\rho$  such that  $|p_n - x_n| \leq \varepsilon_1$ . Thus  $x_n \in K^{\varepsilon_1}$  for all  $n \in \mathbb{N}$ , and  $x_0 \in \text{Viab}_{\Gamma_\rho}(K^{\varepsilon_1})$  by Observation 1. □

The following lemma uses a simple calculation: For any  $A \subset X$ , we have  $\text{dist}(A, A_h^{\alpha(h)}) \leq \alpha(h)$ , because for every  $a \in A$  there is an  $x_h \in X_h$  such that  $|a - x_h| \leq \alpha(h)$ , and then  $x_h \in A^{\alpha(h)} \cap X_h$ .

**Lemma 6.** *If  $\varepsilon_1 + \alpha(h) \leq \varepsilon_0$ , then*

$$\text{dist}(\text{Viab}_{\Gamma_\rho}(K^{\varepsilon_1}), \text{Viab}_{\Gamma_{\rho,h}}(K_h^{\varepsilon_1 + \alpha(h)})) \leq \alpha(h).$$

*Proof.* Let  $(x_n)_{n \in \mathbb{N}}$  be a viable orbit of  $\Gamma_\rho$  in  $K^{\varepsilon_1}$ . By definition, there exists a  $\xi_0 \in K_h^{\varepsilon_1 + \alpha(h)}$  such that  $|x_0 - \xi_0| \leq \alpha(h)$ . Since

$$\text{dist}(x_0 + \rho F(x_0), \xi_0 + \rho F(\xi_0)) \leq (1 + L\rho)\alpha(h), \quad (10)$$

we have

$$\text{dist}(x_0 + \rho F(x_0), \xi_0 + \rho F(\xi_0) + (1 + L\rho)\alpha(h)B) = 0, \quad (11)$$

and thus

$$\begin{aligned} & \text{dist}(\Gamma_\rho(x_0), \Gamma_{\rho,h}(\xi_0)) \\ &= \text{dist}(x_0 + \rho F(x_0), (\xi_0 + \rho F(\xi_0) + (2 + L\rho)\alpha(h)B) \cap X_h) \\ &\leq \alpha(h). \end{aligned}$$

Thus there exists a  $\xi_1 \in \Gamma_{\rho,h}(\xi_0)$  such that  $|x_1 - \xi_1| \leq \alpha(h)$ , and by induction there exists a whole orbit  $(\xi_n)_{n \in \mathbb{N}}$  of  $\Gamma_{\rho,h}$  with  $|x_n - \xi_n| \leq \alpha(h)$  for all  $n \in \mathbb{N}$ . Consequently  $\xi_n \in K_h^{\varepsilon_1 + \alpha(h)}$  for all  $n \in \mathbb{N}$ , and  $\xi_0 \in \text{Viab}_{\Gamma_{\rho,h}}(K_h^{\varepsilon_1 + \alpha(h)})$ .  $\square$

**Lemma 7.** *Let  $\varepsilon_2 := \varphi(k(\rho, h) + M\rho(e^{L\rho} - 1))$ . If  $M\rho e^{L\rho} \leq \varepsilon_0 - \varepsilon_1 - \alpha(h)$ ,  $M\rho \leq \varepsilon_0 - \varepsilon_1 - \alpha(h) - \varepsilon_2$ , and  $k(\rho, h) + M\rho(e^{L\rho} - 1) \leq d_0^{(s)}$ ,*

$$\text{dist}(\text{Viab}_{\Gamma_{\rho,h}}(K_h^{\varepsilon_1 + \alpha(h)}), \text{Viab}_F(K)) \leq \varepsilon_2.$$

*Proof.* By Lemma 4,

$$\begin{aligned} & \text{dist}(\Gamma_{\rho,h}(x_h), G_\rho(x_h)) \\ &\leq \text{dist}(x_h + \rho F(x_h) + k(\rho, h)B, x_h + \rho F(x_h)) + \text{dist}(x_h + \rho F(x_h), G_\rho(x_h)) \\ &\leq k(\rho, h) + M\rho(e^{L\rho} - 1) =: d \end{aligned}$$

for every  $x_h \in K_h^{\varepsilon_1 + \alpha(h)}$ . Thus any trajectory  $(\xi_n)_{n \in \mathbb{N}}$  of  $\Gamma_{\rho,h}$  which is viable in  $K_h^{\varepsilon_1 + \alpha(h)}$  is a  $d$ -pseudotrajectory of  $G_\rho$ , and assumption (iia) implies the existence of an orbit  $(p_n)_{n \in \mathbb{N}}$  of  $G_\rho$  such that  $|p_n - \xi_n| \leq \varepsilon_2$  for all  $n \in \mathbb{N}$ . Hence  $x_0 \in \text{Viab}_{G_\rho}(K^{\varepsilon_1 + \alpha(h) + \varepsilon_2}) = \text{Viab}_F(K)$  by Lemma 3 and Observation 1.  $\square$

Altogether we obtain an estimate for the accuracy of the viability kernel algorithm:

**Theorem 8.** *If*

$$k(\rho, h) + M\rho(e^{L\rho} - 1) \leq d_0^{(s)}, \quad (12)$$

$$M\rho(e^{L\rho} - 1) \leq d_0^{(is)}, \quad (13)$$

$$M\rho e^{L\rho} + \varepsilon_1 + \alpha(h) \leq \varepsilon_0, \quad (14)$$

$$M\rho + \varepsilon_1 + \varepsilon_2 + \alpha(h) \leq \varepsilon_0, \quad (15)$$

*and if assumptions (i) to (iii) are satisfied, then*

$$\text{dist}_H(\text{Viab}_F(K), \text{Viab}_{\Gamma_{\rho,h}}(K_h^{\varepsilon_1 + \alpha(h)})) \leq \max\{\varepsilon_1 + \alpha(h), \varepsilon_2\}. \quad (16)$$

The conditions of Theorem 8 do not look very appealing. Please note that they can be verified easily in a quite practical sense: Let a desired accuracy  $\delta > 0$  of the approximation of the viability kernel be prescribed. If concrete monotonous functions  $\varphi$  and  $\psi$  are given, the inequalities (12) to (15) together with

$$\max\{\varepsilon_1 + \alpha(h), \varepsilon_2\} \leq \delta \quad (17)$$

can be regarded as constraints which are monotonous w.r.t.  $\rho$  and  $\alpha(h)$ . One can fix a (sufficiently small)  $\rho > 0$  and determine the maximal  $\alpha(h)$  such that the pair  $(\rho, \alpha(h))$  satisfies all constraints using a simple interval subdivision algorithm. The same method can be used in the context of Theorem 11.

### 3.3 Estimates using the shadowing property only

It is possible to dispense with the inverse shadowing property by inflating the right hand sides of the numerical schemes so much that the numerical errors are 'swallowed' by the inflation. Therefore, we define

$$\tilde{\Gamma}_\rho : X \rightrightarrows X, \quad x \mapsto x + \rho F(x) + M\rho(e^{L\rho} - 1)B \quad (18)$$

and

$$\tilde{\Gamma}_{\rho,h} : X_h \rightrightarrows X_h, \quad x_h \mapsto (x_h + \rho F(x_h) + M\rho(e^{L\rho} - 1)B + k(\rho, h)B) \cap X_h. \quad (19)$$

For these schemes we have

**Lemma 9.** *If  $M\rho e^{L\rho} \leq \varepsilon_0$  and  $\alpha(h) \leq \varepsilon_0$ ,*

$$\text{dist}(\text{Viab}_F(K), \text{Viab}_{\tilde{\Gamma}_{\rho,h}}(K_h^{\alpha(h)})) \leq \alpha(h).$$

*Proof.* According to Lemma 4,

$$\text{dist}(G_\rho(x_0), \Gamma_\rho(x_0)) \leq M\rho(e^{L\rho} - 1)$$

for all  $x_0 \in K$ , and thus

$$\text{dist}(G_\rho(x_0), \tilde{\Gamma}_\rho(x_0)) = 0$$

and

$$\text{dist}(\text{Viab}_{G_\rho}(K), \text{Viab}_{\tilde{\Gamma}_\rho}(K)) = 0.$$

Adapting the proof of Lemma 6 to  $\tilde{\Gamma}_\rho$  and  $\tilde{\Gamma}_{\rho,h}$  yields the desired result.  $\square$



**Lemma 10.** *If  $\varepsilon_3 := \varphi(2M\rho(e^{L\rho} - 1) + k(\rho, h)) \leq \varepsilon_0 - \alpha(h)$ ,  $M\rho e^{L\rho} \leq \varepsilon_0 - \alpha(h)$ , and  $2M\rho(e^{L\rho} - 1) + k(\rho, h) \leq d_0^{(s)}$ , then*

$$\text{dist}(\text{Viab}_{\tilde{\Gamma}_{\rho,h}}(K_h^{\alpha(h)}), \text{Viab}_F(K)) \leq \varepsilon_3.$$

*Proof.* By Lemma 4,

$$\begin{aligned} & \text{dist}(\tilde{\Gamma}_{\rho,h}(x_h), G_\rho(x_h)) \\ & \leq \text{dist}(x_h + \rho F(x_h) + M\rho(e^{L\rho} - 1)B + k(\rho, h)B, x_h + \rho F(x_h)) \\ & \quad + \text{dist}(x_h + \rho F(x_h), G_\rho(x_h)) \\ & \leq 2M\rho(e^{L\rho} - 1) + k(\rho, h) =: \tilde{d} \end{aligned}$$

for any  $x_h \in K_h^{\alpha(h)}$ .

Thus any trajectory  $(\xi_n)_{n \in \mathbb{N}}$  of  $\tilde{\Gamma}_{\rho,h}$  which is viable in  $K_h^{\alpha(h)}$  is a  $\tilde{d}$ -pseudotrajectory of  $G_\rho$ , and assumption (ia) implies that there exists an orbit  $(p_n)_{n \in \mathbb{N}}$  of  $G_\rho$  such that  $|p_n - \xi_n| \leq \varepsilon_3$  for all  $n \in \mathbb{N}$ . Hence  $p_0 \in \text{Viab}_{G_\rho}(K^{\alpha(h)+\varepsilon_3}) = \text{Viab}_F(K)$  by Lemma 3.  $\square$

Summarizing we obtain an estimate for the accuracy of the viability kernel algorithm for systems which have the shadowing but not the inverse shadowing property:

**Theorem 11.** *If*

$$M\rho e^{L\rho} + \alpha(h) \leq \varepsilon_0, \quad (20)$$

$$2M\rho(e^{L\rho} - 1) + k(\rho, h) \leq d_0^{(s)}, \quad (21)$$

$$\varepsilon_3 + \alpha(h) \leq \varepsilon_0, \quad (22)$$

*and if assumptions (i), (ia), and (iii) are satisfied, then*

$$\text{dist}_H(\text{Viab}_F(K), \text{Viab}_{\tilde{\Gamma}_{\rho,h}}(K_h^{\alpha(h)})) \leq \max\{\alpha(h), \varepsilon_3\}. \quad (23)$$

## 4 Example

It was shown in [9] that certain differential inclusions with a contractive  $\rho$ -flow have the shadowing and the inverse shadowing property. The results which are relevant for this paper are summarized in the following

**Theorem 12.** *Let the right hand side  $F : X \rightrightarrows X$  of (1) be Lipschitz continuous with constant  $L$  and with compact and convex values, and let  $\|F(x)\| \leq M$  for some  $M > 0$  and all  $x \in X$ . Assume that there exist*

numbers  $a > 0$  and  $\lambda \in (0, 1)$  such that the  $\rho$ -flow defined by (2) satisfies the stability condition

$$\text{dist}_H(G_\rho(x), G_\rho(x + v)) \leq \lambda|v| \quad (24)$$

for any  $x \in X$  and  $|v| \leq a$ . If  $\rho \leq \frac{d}{2M}$  and  $d < d_0 := \frac{1-\lambda}{2}a$ , then  $G_\rho$  has the  $(d, \frac{2d}{1-\lambda})$ -shadowing property and the inverse  $(d, \frac{2d}{1-\lambda})$ -shadowing property in  $X$ .

In the same paper it was shown that a differential inclusion satisfies (24) if its right hand side has the relaxed one-sided Lipschitz property with a negative Lipschitz constant.

Theorem 12 is formulated in such a way that all assumptions are required to be true on the whole space. However, it is also valid if the assumptions are satisfied on a subset  $K^{\varepsilon_0}$ ,

- the given  $d$ -pseudotrajectory of  $G_\rho$  is contained in  $K^\varepsilon$  (shadowing)
- the given orbit of  $G_\rho$  is contained in  $K^\varepsilon$  (inverse shadowing)

for some  $\varepsilon > 0$ , and  $a \leq \varepsilon_0 - \varepsilon$ , because the behaviour of  $G_\rho$  is only relevant on an  $a$ -neighbourhood of the given trajectory.

Repeating the line of argument of section 3.2 in this setup, we obtain

**Theorem 13.** *Let the right hand side  $F : X \rightrightarrows X$  of (1) be Lipschitz continuous in  $K^{\varepsilon_0}$  with constant  $L$  and with compact and convex values, and let  $\|F(x)\| \leq M$  for some  $M > 0$  and all  $x \in K^{\varepsilon_0}$ . Let  $\lambda \in (0, 1)$  such that the  $\rho$ -flow  $G_\rho$  satisfies (24) for any  $x \in K^{\varepsilon_0}$  and  $v \in X$  with  $x + v \in K^{\varepsilon_0}$ . If*

$$\frac{5-\lambda}{1-\lambda}\alpha(h) + \frac{2L}{1-\lambda}\alpha(h)\rho + \frac{4}{1-\lambda}M\rho(e^{L\rho} - \frac{3+\lambda}{4}) \leq \varepsilon_0$$

then

$$\text{dist}_H(\text{Viab}_F(K), \text{Viab}_{\Gamma_{\rho,h}}(K_h^{\varepsilon_1 + \alpha(h)})) \leq \frac{4}{1-\lambda}\alpha(h) + \frac{2L}{1-\lambda}\alpha(h)\rho + \varepsilon_1,$$

where  $\varepsilon_1 = \frac{2}{1-\lambda}M\rho(e^{L\rho} - 1)$ .

Following section 3.3, we obtain a slightly but not substantially worse estimate under similar conditions.

In classical numerical analysis, the order of convergence of a scheme is regarded as one of the most important indicators for its quality. It is doubtful if this way of thinking is appropriate here, but it is possible to obtain linear convergence w.r.t. the spatial discretization  $\alpha(h)$  by setting  $\alpha(h) := \rho$ .

Please note that every shadowing theorem for the  $\rho$ -flow of a differential inclusion can be used to derive a concrete error estimate for the viability kernel algorithm in the way sketched above. As the shadowing theory for set-valued systems is still being developed, there is hope that the reasoning of section 3 can soon be applied to more general classes of right hand sides.

## References

- [1] E. Akin. *Simplicial Dynamical Systems*. volume 667 of Memoires of the Amer. Math. Soc., 1999.
- [2] J.-P. Aubin. *Viability Theory*. Birkhäuser, Boston, 1991.
- [3] J.P. Aubin and A. Cellina. *Differential Inclusions*, volume 264 of *Grundlehren der mathematischen Wissenschaften*. Springer, Berlin, 1984.
- [4] E. Crück and P. Saint-Pierre. Nonlinear impulse target problems under state constraint: A numerical analysis based on viability theory. *Set-Valued Analysis*, 12:383–416, 2004.
- [5] H. Frankowska and M. Quincampoix. Viability kernels of differential inclusions with constraints: Algorithm and applications. *J. Math. Systems Estimation Control*, 1(3):371–388, 1991.
- [6] Vasile Glavan and Valeriu Gutu. On the dynamics of contracting relations. *Analysis and optimization of differential systems*, Kluwer Acad. Publ., Boston, MA, pages 179–188, 2003.
- [7] K. Palmer. *Shadowing in Dynamical Systems. Theory and Applications*. Kluwer, Dordrecht, 2000.
- [8] Sergei Yu. Pilyugin. *Shadowing in Dynamical Systems*, volume 1706 of *Lect. Notes Math*. Springer, Berlin, 1999.
- [9] Sergei Yu. Pilyugin and Janosch Rieger. Shadowing and inverse shadowing in set-valued dynamical systems. contractive case. *Top. Meth. Nonl. Anal.*, accepted in 2007.
- [10] Sergei Yu. Pilyugin and Janosch Rieger. Shadowing and inverse shadowing in set-valued dynamical systems. hyperbolic case. *Top. Meth. Nonl. Anal.*, submitted in 2007.

- [11] P. Saint-Pierre. Approximation of the viability kernel. *Appl. Math. Optim.*, 29:187–209, 1994.
- [12] Evelyn Sander. Hyperbolic sets for noninvertible maps and relations. *Disc. Cont. Dyn. Syst.*, 8(2):339–357, 1999.