

FLEXIBILITY OF SURFACE GROUPS IN CLASSICAL SIMPLE LIE GROUPS

INKANG KIM AND PIERRE PANSU

ABSTRACT. We show that a surface group of high genus contained in a classical simple Lie group can be deformed to become Zariski dense, unless the Lie group is $SU(p, q)$ (resp. $SO^*(2n)$, n odd) and the surface group is maximal in some $S(U(p, p) \times U(q - p)) \subset SU(p, q)$ (resp. $SO^*(2n - 2) \times SO(2) \subset SO^*(n)$). This is a converse, for classical groups, to a rigidity result of S. Bradlow, O. García-Prada and P. Gothen.

Dedicated to Lionel Bérard-Bergery, for his 65th birthday

1. INTRODUCTION

Free groups are obviously flexible. In particular, a generic free subgroup in a real algebraic group is Zariski dense. This already fails for surface groups, although they are very flexible from other points of view. The first evidence came from the following result by D. Toledo.

Theorem 1.1. (D. Toledo, 1979, 1989, [15]). *Let Γ be a discrete cocompact subgroup of $SU(1, 1)$. Map $SU(1, 1)$ as a 2×2 block in $SU(1, n)$, $n \geq 2$. Then every neighboring homomorphism $\Gamma \rightarrow SU(1, n)$ is contained in a conjugate of $S(U(1, 1) \times U(n - 1))$.*

In fact, Toledo obtained a stronger, global result: a characterization of surface subgroups of $S(U(1, 1) \times U(n - 1))$ among surface subgroups of $SU(1, n)$ by the value of a characteristic class known as Toledo's invariant, which we now define.

Let X be a Hermitian symmetric space, with Kähler form Ω (the metric is normalized so that the minimal sectional curvature equals -1). Let Σ be a closed surface of negative Euler characteristic, let $\Gamma = \pi_1(\Sigma)$ act isometrically on X . Pick a smooth equivariant map $\tilde{f} : \tilde{\Sigma} \rightarrow X$.

Definition 1.2. *Define the Toledo invariant of the action $\rho : \Gamma \rightarrow \text{Isom}(X)$ by*

$$T_\rho = \frac{1}{2\pi} \int_\Sigma \tilde{f}^* \Omega.$$

Then

- (1) T_ρ depends continuously on ρ .

¹I. Kim gratefully acknowledges the partial support of KRF Grant (0409-20060066) and a warm support of IHES during his stay.

- (2) There exists $\ell_X \in \mathbb{Q}$ such that $T_\rho \in \ell_X \mathbb{Z}$.
- (3) $|T_\rho| \leq |\chi(\Sigma)| \text{rank}(X)$.

Definition 1.3. *Actions ρ such that $|T_\rho| = |\chi(\Sigma)| \text{rank}(X)$ are called maximal representations.*

The following result generalizes Theorem 1.1.

Theorem 1.4. (L. Hernández Lamóneda, [11], S. Bradlow, O. García-Prada, P. Gothen, [1], [2]). *Maximal reductive representations of Γ to $SU(p, q)$, $p \leq q$, can be conjugated into $S(U(p, p) \times U(q - p))$.*

In turn, Theorem 1.4 is a special case of a more general result.

Definition 1.5. *Say a Hermitian symmetric space is of tube type if it can be realized as a domain in \mathbb{C}^n of the form $\mathbb{R}^n + iC$ where $C \subset \mathbb{R}^n$ is a proper open cone.*

Example 1.6. *Siegel's upper half spaces and Grassmannians with isometry groups $PO(2, q)$ are of tube type.*

The Grassmannian $\mathcal{D}_{p, q}$, $p \leq q$, with isometry group $PU(p, q)$ is of tube type iff $p = q$.

The Grassmannian \mathcal{G}_n with isometry group $SO^(2n)$ is of tube type iff n is even.*

The exceptional Hermitian symmetric space of dimension 27 is of tube type, the other one (of dimension 16) is not.

Products of tube type spaces are of tube type, so polydisks are of tube type.

Remark 1.7. *All maximal tube type subsymmetric spaces in a Hermitian symmetric space are conjugate. For instance, the maximal tube type subsymmetric space in $\mathcal{D}_{p, q}$ is $\mathcal{D}_{p, p}$. The maximal tube type subsymmetric space in \mathcal{G}_{2n+1} is \mathcal{G}_{2n} .*

Theorem 1.8. (Burger, Iozzi, Wienhard, [3]). *Let Γ be a closed surface group and X a Hermitian symmetric space. Every maximal representation $\Gamma \rightarrow \text{Isom}(X)$ stabilizes a maximal tube type subsymmetric space Y . Conversely, for every tube type Hermitian symmetric space X , $\text{Isom}(X)$ admits Zariski dense maximal surface subgroups.*

1.1. Results. Our main result is a converse of Theorem 1.4 (i.e. Theorem 1.8 for classical simple Lie groups).

Theorem 1. *Let G be a classical real Lie group, i.e. a real form of $SL(n, \mathbb{C})$, $O(n, \mathbb{C})$ or $Sp(n, \mathbb{C})$. Let Γ be the fundamental group of a closed surface of genus $\geq 2 \dim(G)^2$. A homomorphism $\phi : \Gamma \rightarrow G$ can be approximated by Zariski dense representations, unless the symmetric space X of G is Hermitian and not of tube type, and ϕ is maximal.*

In other words, the exceptions are $G = SU(p, q)$, $q > p$ and $\phi(\Gamma)$ is contained in a conjugate of $S(U(p, p) \times U(q - p)) \subset SU(p, q)$, or $G = SO^*(2n)$, n odd, and $\phi(\Gamma)$ is contained in a conjugate of $SO^*(2n - 2) \times SO(2) \subset SO^*(n)$.

The genus assumption is probably unnecessary, but we are unable to remove it.

Question 1.9. *Does Theorem 1 extend to exceptional simple Lie groups ?*

Since flexibility easily holds in compact and complex groups (see Proposition 4.3), and the rank one example F_4^{-20} is treated in [12], there remains 10 cases ($E_6^6, E_6^{-14}, E_6^{-26}, E_7^7, E_7^{-5}, E_7^{-25}, E_8^8, E_8^{-24}, F_4^4, G_2^2$) with at least one rigidity case (E_6^{-14}).

1.2. Scheme of proof. The proof relies on

- a necessary and sufficient condition for flexibility from [12], see Theorem 2.2 below;
- tools from Burger, Iozzi and Wienhard's theory of tight maps between Hermitian symmetric spaces, [4];
- a detailed analysis of centers of centralizers of reductive subgroups of classical simple Lie groups.

This last analysis is performed in a rather bare handed manner, based on bilinear and sesquilinear algebra. This is where exceptional simple Lie groups elude us.

1.3. Plan of the paper. Section 2 recalls the needed result from [12]. Section 3 proves relevant consequences of the theory of tight maps. Section 4 provides a description of classical real simple Lie groups as fixed points of involutions which helps in computing root space decompositions in the complexified Lie algebra. This is done in section 5 for $\mathfrak{sl}(n, \mathbb{C})$ and in section 6 for $\mathfrak{so}(n, \mathbb{C})$ and $\mathfrak{sp}(n, \mathbb{C})$. The method consists in first computing the root space decomposition in the standard representation of the complexified Lie algebra, and deducing the decomposition in the adjoint representation. The theory of tight homomorphisms allows to exclude balancedness (with exceptions), first for real forms of $SL(n, \mathbb{C})$ in section 7, then for real forms of $SO(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$ in section 8. Theorem 1 is proven in section 9.

2. FLEXIBILITY CRITERION

As far as the flexibility of a homomorphism ϕ is concerned, a key role is played by the center of the centralizer of the image of ϕ . It splits the complexified Lie algebra of G into root spaces \mathfrak{g}_λ . When the root λ is pure imaginary, \mathfrak{g}_λ carries a natural nondegenerate sesquilinear form defined as follows. Let $(X, X') \mapsto X \cdot X'$ denote the Killing form on $\mathfrak{g} \otimes \mathbb{C}$. Then the sesquilinear form

$$s_\lambda(X, X') = \bar{X} \cdot X'$$

is nondegenerate on \mathfrak{g}_λ . If $p - q$ is the signature of s_λ , the representation of Γ on \mathfrak{g}_λ gives rise to a homomorphism $\Gamma \rightarrow U(\mathfrak{g}_\lambda, s_\lambda) = U(p, q)$ and an

isometric action on the Grassmannian $\mathcal{D}_{p,q}$, and thus a Toledo invariant T_λ (by convention, $\mathcal{D}_{0,q}$ is a point and T_λ vanishes if $p = 0$)¹.

Definition 2.1. *Let $\mathfrak{t} \subset \mathfrak{g}$ be a torus, centralized by a homomorphism $\phi : \Gamma \rightarrow G$. Among the roots of the adjoint action of \mathfrak{t} on \mathfrak{g} , let P be the subset of pure imaginary roots λ such that $2T_\lambda = -\chi(\Gamma)\dim(\mathfrak{g}_\lambda)$. Say \mathfrak{t} is balanced with respect to ϕ if 0 belongs to the interior of the sum of the convex hull of the imaginary parts of elements of P and the linear span of the real and imaginary parts of roots not in $\pm P$.*

Here is a necessary and sufficient condition for flexibility, for surface groups of sufficiently large genus.

Theorem 2.2. ([12], Theorem 3). *Let G be a semisimple real algebraic group. Let Γ be the fundamental group of a closed surface of genus $\geq 2\dim(G)^2$. Let $\phi : \Gamma \rightarrow G$ be a homomorphism with reductive Zariski closure. Then ϕ is flexible if and only if \mathfrak{c} , the center of the centralizer of $\phi(\Gamma)$, is balanced with respect to ϕ .*

The proof of Theorem 1 will rely on this criterion: we shall describe centers of centralizers of reductive subgroups and their roots and pile up restrictions on the set P that make non balancedness exceptional. The reduction from nonreductive to reductive homomorphisms will be explained in section 9.

3. TIGHTNESS

Definition 3.1. (Burger, Iozzi, Wienhard, [4]). *Let G be a reductive Hermitian group, i.e. a connected reductive Lie group in which the center is compact and such that the symmetric spaces X_i associated to all simple noncompact factors is Hermitian. Normalize the metric on X_i so that the minimum holomorphic sectional curvature equals -1 . Let κ_G^b denote the bounded continuous cohomology class on G defined by integrating the Kähler form of $X = \coprod X_i$ on triangles with geodesic sides. Let Γ be locally compact group. Say a continuous homomorphism $\phi : \Gamma \rightarrow G$ is tight if*

$$\|\phi^* \kappa_G^b\| = \|\kappa_G^b\|.$$

Proposition 3.2. *Let G be a semisimple real algebraic group. Let Γ be a surface group. Let $\phi : \Gamma \rightarrow G$ be a homomorphism. Let \mathfrak{c} denote the center of the centralizer of $\phi(\Gamma)$. Let $Z_G(\mathfrak{c})$ denote its centralizer in G . Let λ be a pure imaginary root of the adjoint action of \mathfrak{c} on $\mathfrak{g} \otimes \mathbb{C}$. Let \mathfrak{g}_λ denote the corresponding root space and T_λ the Toledo invariant corresponding to the natural sesquilinear form s_λ on \mathfrak{g}_λ . Assume that*

$$2T_\lambda = -\chi(\Gamma)\dim(\mathfrak{g}_\lambda).$$

Then

¹This definition is equivalent to Definition 1.15 and Corollary 4.8 of [12]. Indeed, the symplectic structure on a real version of \mathfrak{g}_λ alluded to there is the imaginary part of s_λ .

- (1) s_λ has vanishing signature;
- (2) $\rho_\lambda : \Gamma \rightarrow U(g_\lambda, s_\lambda)$ is maximal;
- (3) the homomorphism $Z_G(\mathfrak{c}) \rightarrow U(g_\lambda, s_\lambda)$ is tight.

Proof: 1. Let $\sigma = p - q$ be the signature of the sesquilinear form s_λ , with $p + q = \dim_{\mathbb{C}}(\mathfrak{g}_\lambda)$. By assumption,

$$2T_\lambda = -\chi(\Gamma)(p + q).$$

On the other hand, the Milnor-Wood inequality for actions on Hermitian symmetric spaces (due to J. Milnor, [13], V. Turaev, [16], A. Domic and D. Toledo, [6], J.-L. Clerc and B. Ørsted, [5]), states that

$$(1) \quad |T_\lambda| \leq |\chi(\Gamma)| \text{rank}(\mathcal{D}_{p,q}) = |\chi(\Gamma)| \min\{p, q\}.$$

Thus $p + q \leq 2 \min\{p, q\}$. This implies that $p = q$.

2. Also, inequality (1) has to be an equality, so $\rho_\lambda : \Gamma \rightarrow U(g_\lambda, s_\lambda)$ is maximal.

3. A maximal representation of a surface group to a semisimple group of Hermitian type is a special case of a tight homomorphism. Since ρ_λ factors via the homomorphism $Z_G(\mathfrak{c}) \rightarrow U(g_\lambda, s_\lambda)$, this has to be tight itself. ■

The following two facts will play a key role in our argument.

Lemma 3.3. *Let $\rho : G \rightarrow G'$ be a tight homomorphism between semisimple groups of Hermitian type. If the kernel of ρ is compact and G' is of tube type, so is G .*

Proof: This is Theorem 6.2 of [4]. ■

Lemma 3.4. *Let V, V' be vector spaces equipped with nondegenerate sesquilinear forms. Assume that $\text{Hom}(V, V')$, equipped with the natural sesquilinear form*

$$(f, f') \mapsto \text{Trace}(f^* \circ f'),$$

(here, f^ denotes the adjoint with respect to the sesquilinear forms on V and V'), has vanishing signature. Assume that the induced homomorphism $U(V) \times U(V') \rightarrow U(\text{Hom}(V, V'))$ is tight (see Definition 3.1). Then one of V and V' is definite and the other has vanishing signature.*

Proof: Since $U(\text{Hom}(V, V'))$ is Hermitian of tube type, Lemma 3.3 implies that $U(V) \times U(V')$ is of tube type, up to compact groups. This implies that each of the sesquilinear vector spaces V and V' is either of vanishing signature or definite. Clearly, if both are definite, $U(V) \times U(V')$ is compact and tightness is out of sight.

Let us show that V and V' cannot both have vanishing signature, i.e. one of them must be definite. For this, we use the tightness criterion of [4], Corollary 8.2. Let J_ℓ denote the generator of the center of the maximal compact subgroup of $U(V)$ which defines the complex structure on the symmetric space of $U(V)$. In a splitting of $V = V^+ \oplus V^-$ in a sum of positive

(resp. negative) definite subspaces,

$$J_V = \begin{pmatrix} \frac{i}{2}I_{d_V} & 0 \\ 0 & -\frac{i}{2}I_{d_V} \end{pmatrix}$$

Use a similar splitting $V' = V'^+ \oplus V'^-$ and split accordingly $Hom(V, V')$, whose elements become block matrices $N = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. The action of

$U_V \times U_{V'}$ is $(U, U', N) \mapsto U'^{-1}NU$. The linearized action is $(U, U', N) \mapsto -U'N + NU$. The image of $(J_V, J_{V'})$ under the linearized action is $N \mapsto$

$$\begin{pmatrix} 0 & -iB \\ iC & 0 \end{pmatrix}, \text{ a map whose matrix is } \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \end{pmatrix} \text{ in a splitting of } Hom(V, V')$$

into a positive subspace $Hom(V^+, V'^+) \oplus Hom(V^-, V'^-)$ and a negative subspace $Hom(V^+, V'^-) \oplus Hom(V^-, V'^+)$. Therefore the inner product with the element

$$J_{Hom(V, V')} = \begin{pmatrix} \frac{i}{2}I_{d_V} & 0 & 0 & 0 \\ 0 & \frac{i}{2}I_{d_V} & 0 & 0 \\ 0 & 0 & -\frac{i}{2}I_{d_V} & 0 \\ 0 & 0 & 0 & -\frac{i}{2}I_{d_V} \end{pmatrix}$$

vanishes. According to [4], Corollary 8.2, the homomorphism $U(V) \times U(V') \rightarrow U(Hom(V, V'))$ is not tight. We conclude that one of V and V' has vanishing signature and the other one is definite. ■

Lemma 3.5. *Let Γ be a surface group. Let U be a unitary 1-dimensional representation of Γ and W a sesquilinear representation of Γ . If $Hom(U, W)$ is a maximal representation if and only if W is.*

Proof: $Hom(I_0, W)$ and W are isomorphic as projective representations. Since $U(W)$ acts on the symmetric space of $SU(W)$ via its quotient $PU(W)$, maximality is a projectively invariant property. ■

Lemma 3.6. *Let Γ be a surface group. Let W, W' be sesquilinear representations of Γ . Then*

- (1) $T(\bar{W}) = -T(W)$.
- (2) $T(W \oplus W') = T(W) + T(W')$.

Proof: 1. Passing from W to \bar{W} changes the sign of the complex structure on the symmetric space X of $SU(W)$. This changes the sign of the Kähler form, and thus the sign of Toledo invariants.

2. When $X \times X'$ is mapped to the symmetric space Y of $SU(W \oplus W')$, the Kähler form Ω_Y of Y restricts on the complex totally geodesic manifold $X \times X'$ to $\Omega_X + \Omega_{X'}$, so Toledo invariants add up. ■

Lemma 3.7. *The injection $\iota : O(2, 2) \hookrightarrow U(2, 2)$ is not tight.*

Proof: $O(2, 2) \subset U(2, 2)$ is the fixed point set of conjugation σ , i.e. $\sigma \circ \iota = \sigma$. σ induces an orientation reversing isometry of the symmetric space X of $U(2, 2)$, which changes the sign of the Kähler form, $\sigma^*\Omega_X = -\Omega_X$. ι induces a totally geodesic embedding of symmetric spaces, still denoted by $\iota : Y \rightarrow X$. Since $\iota^*\Omega_X = \iota^*\sigma^*\Omega_X = -\iota^*\Omega_X$, $\iota^*\Omega_X = 0$, ι^* is not isometric in bounded continuous cohomology, so ι is not tight. ■

Lemma 3.8. *Let Γ be a surface group. Let $\iota : Y \rightarrow X$ be a holomorphic and isometric embedding of Hermitian symmetric spaces of equal rank. Let $\rho : H \rightarrow G$ denote the corresponding homomorphism between isometry groups. Let $\phi : \Gamma \rightarrow H$ be a homomorphism. If $\rho \circ \phi$ is a maximal representation, so is ϕ .*

Proof: By assumption, Kähler forms match, i.e. $\iota^*\Omega_X = \Omega_Y$, thus Toledo invariants coincide, $T(\phi) = T(\rho \circ \phi)$. Since X and Y have the same rank, $\rho \circ \phi$ is maximal if and only if ϕ is. ■

4. CLASSICAL SIMPLE LIE GROUPS

4.1. Definition. Classical simple real Lie groups are special linear groups of division rings D with center \mathbb{R} and special unitary groups of nondegenerate binary forms over D , see the appendix in [14].

There are only 3 such division rings: \mathbb{R} , \mathbb{C} and \mathbb{H} , leading to 3 special linear groups, $SL(n, \mathbb{R})$, $SL(n, \mathbb{C})$, and $SL(n, \mathbb{H})$.

Let ι be a continuous (anti)-automorphism of D : ι can be identity, complex conjugation or quaternionic conjugation. Let $\epsilon \in \pm 1$. By a (ι, ϵ) -symmetric binary² form on a right D -vectorspace V , we mean a D -valued \mathbb{R} -bilinear map $H : V \times V \rightarrow D$ such that

- (1) for all $v, v' \in V$ and all $q \in D$, $H(v, v'q) = H(v, v')q$;
- (2) for all $v, v' \in V$, $H(v', v) = \epsilon\iota(H(v, v'))$.

Say H is *nondegenerate* if the only $v \in V$ such that $H(v, v') = 0$ for all $v' \in V$ is 0.

The group $U(V, H)$ consists of right D -linear self maps of V which preserve H . We are interested in the special unitary group $SU(V, H) = U(V, H) \cap SL(V, D)$.

When $D = \mathbb{R}$, the only choice for ι is identity, leading to

- if $\epsilon = 1$, real special orthogonal groups, indexed by dimension and signature and denoted by $SO(p, q)$, where $p + q = \dim_{\mathbb{R}}(V)$;
- if $\epsilon = -1$, real symplectic groups, indexed by dimension only and denoted by $Sp(n, \mathbb{R})$, where $n = \dim_{\mathbb{R}}(V)$ is even.

When $D = \mathbb{C}$ there are two choice for ι . If ι is identity, this leads to complex special orthogonal (if $\epsilon = 1$) and symplectic (if $\epsilon = -1$) groups, indexed by dimension and denoted respectively by $SO(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$.

²We keep the words Hermitian for positive definite forms, and sesquilinear for the complex case.

If ι is complex conjugation, $\epsilon = \pm 1$ lead to isomorphic groups called special unitary groups, indexed by dimension and signature and denoted by $SU(p, q)$, where $p + q = \dim_{\mathbb{C}}(V)$.

When $D = \mathbb{H}$ there are two choices for ι . If ι is identity, no nonzero (ι, ϵ) -symmetric forms exist. If ι is quaternionic conjugation, this leads to

- if $\epsilon = 1$, quaternionic unitary groups, indexed by dimension and signature and denoted by $Sp(p, q)$, $p + q = \dim_{\mathbb{H}}(V)$;
- if $\epsilon = -1$, quaternionic skew-unitary groups, indexed by dimension only and denoted by $SO^*(2n)$, where $n = \dim_{\mathbb{H}}(V)$ is even.

4.2. Real forms of complex groups. We shall be mainly concerned with the 7 families of non complex groups, 3 attached to binary forms on real vectorspaces, 1 on complex vectorspaces and 3 on quaternionic vectorspaces. Each of these groups is obtained as the fixed point set of an anti- \mathbb{C} -linear involutive automorphism σ of a complex Lie group, as we now explain.

Consider first the complex unitary family $U(p, q)$. Here, the data is a nondegenerate sesquilinear form s on a complex vectorspace V . For $f \in GL(V)$, let f^* denote the s -adjoint of f , defined by

$$\forall v, v' \in V, \quad s(f(v), v') = s(v, f^*(v')).$$

Then $\sigma(f) = (f^*)^{-1}$ is an anti- \mathbb{C} -linear involutive automorphism of $GL(V)$ and of $SL(V)$. The fixed point set of σ in $SL(V)$ is $SU(V, s)$.

The 6 remaining families admit a common construction.

Given a real vectorspace $V_{\mathbb{R}}$, let $V = V_{\mathbb{R}} \otimes \mathbb{C}$ and let $\tau =$ conjugation. Note that $\tau^{-1} = \tau$. If b is a nondegenerate quadratic or symplectic form on $V_{\mathbb{R}}$, let $B = b \otimes \mathbb{C}$.

Given a right quaternionic vectorspace $V_{\mathbb{H}}$, pick a basis $(1, i, j, k)$ of \mathbb{H} , use right multiplication by i to turn $V_{\mathbb{H}}$ into a complex vectorspace denoted by V . Let τ be right multiplication by j . Note that $\tau^{-1} = -\tau$. For $q = a + jb \in \mathbb{H}$, $a, b \in \mathbb{C}$, denote by $\mathcal{C}(q) = a$. If H is a $(-, \epsilon)$ -symmetric binary form on $V_{\mathbb{H}}$, let, for $v, v' \in V$,

$$B(v, v') = \mathcal{C}(vj, v').$$

Then B is a $(-\epsilon)$ -symmetric \mathbb{C} -bilinear form on V .

In both cases, τ is anti- \mathbb{C} -linear, $\tau^{-1} = \eta\tau$ for some $\eta \in \pm 1$, and

$$B(\tau(v), \tau(v')) = \overline{B(v, v')}.$$

For $f \in \text{End}_{\mathbb{C}}(V)$, set

$$\sigma(f) = \tau \circ f \circ \tau^{-1}.$$

Then σ is an anti- \mathbb{C} -linear involutive automorphism of the algebra $\mathfrak{gl}(V) = \text{End}_{\mathbb{C}}(V)$ and of its subgroups $SL(V)$ and $O(V, B)$. The fixed point set of σ in $SL(V)$ is $SL(V_{\mathbb{R}})$ (resp. $SL(V_{\mathbb{H}})$). The fixed point set of σ in $O(V, B)$ is $U(V_{\mathbb{R}}, b)$ (resp. $U(V_{\mathbb{H}}, H)$). This construction yields 6 of the 7 families of non complex classical groups.

4.3. Consequences for roots. The special form of the involution σ for 6 of the 7 families of noncomplex groups has the following consequence.

Lemma 4.1. *Let G be a real form of $SL(n, \mathbb{C})$ (resp. $O(n, \mathbb{C})$ or $Sp(n, \mathbb{C})$). Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Let ℓ be a root of \mathfrak{c} on \mathbb{C}^n . Let I_ℓ denote the corresponding root space. Then $\bar{\ell}$ is a root and $I_{\bar{\ell}} = \tau(I_\ell)$. Furthermore,*

- (1) *if $G = SL(n, \mathbb{R})$, $O(p, q)$ or $Sp(n, \mathbb{R})$, $I_{\bar{\ell}} = \bar{I}_\ell$;*
- (2) *if $G = SL(n, \mathbb{H})$, $Sp(p, q)$ or $SO^*(n)$, $I_\ell + I_{\bar{\ell}}$ is a quaternionic subspace.*

Proof: Let $f \in \mathfrak{c}$ and $v \in I_\ell$. Then

$$f(\tau(v)) = \tau(f(v)) = \tau(\ell(f)v) = \overline{\ell(v)}\tau(v)$$

showing that $I_{\bar{\ell}} = \tau(I_\ell)$. For the 3 real families, τ is conjugation, thus $I_{\bar{\ell}} = \bar{I}_\ell$. For the 3 quaternionic families, τ is right multiplication by j , so $I_\ell + I_{\bar{\ell}}$ is stable by right multiplication by i and j , i.e. a quaternionic subspace. ■

4.4. Killing form. Here is a formula for the sesquilinear structure appearing in Theorem 2.2, valid in all cases.

Lemma 4.2. *Let $\mathfrak{g}_{\mathbb{C}} \subset \mathfrak{sl}(n, \mathbb{C})$ be a complex Lie subalgebra of $\mathfrak{sl}(n, \mathbb{C})$. Let σ be an anti- \mathbb{C} -linear involutive automorphism of $\mathfrak{g}_{\mathbb{C}}$, with fixed point set \mathfrak{g} . The natural sesquilinear form s on $\mathfrak{g} \otimes \mathbb{C}$ arising from the Killing form of \mathfrak{g} (see section 2) is proportional to*

$$s(X, X') = \text{Trace}(\sigma(X) \circ X').$$

Proof: Since σ is anti- \mathbb{C} -linear, for $f \in \mathfrak{g}$, $\sigma(if) = -i\sigma(f) = -if$. It follows that the -1 -eigenspace of σ in $\mathfrak{g}^{\mathbb{C}}$ is $i\mathfrak{g}$. The map

$$\mathfrak{g}^{\mathbb{C}} \rightarrow \mathfrak{g} \otimes \mathbb{C}, \quad X \mapsto \left(\frac{X + \sigma(X)}{2}, \frac{X - \sigma(X)}{2i} \right)$$

is an isomorphism. It pulls back conjugation on $\mathfrak{g} \otimes \mathbb{C}$ to σ on $\mathfrak{g}^{\mathbb{C}}$, thus it pulls back the sesquilinear form $\bar{X} \cdot X$ of $\mathfrak{g} \otimes \mathbb{C}$ to $\sigma(X) \cdot X$ on $\mathfrak{g}^{\mathbb{C}}$. The Killing form on $\mathfrak{g}^{\mathbb{C}}$ is proportional to the restriction to $\mathfrak{g}^{\mathbb{C}}$ of the Killing form of $\mathfrak{sl}(n, \mathbb{C})$, whence the formula $\text{Trace}(\sigma(X) \circ X)$. ■

4.5. Flexibility of compact and complex Lie groups. For completeness's sake, we recall here the treatment of compact and complex Lie groups from [12].

Proposition 4.3. *Let G be a compact or complex semisimple Lie group. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of the centralizer of a reductive subgroup. Then \mathfrak{c} is balanced.*

Proof: Let $X \in \mathfrak{g} \otimes \mathbb{C}$, $X = f + ig$. Then $s(X, X) = \bar{X} \cdot X = f \cdot f + g \cdot g$. If G is compact, the Killing form is negative definite, so is s . For all roots λ of \mathfrak{c} in the adjoint representation, the sesquilinear form s_λ on the root

space \mathfrak{g}_λ is negative definite. No root has vanishing signature. According to Proposition 3.2, P is empty so \mathfrak{c} is balanced.

If G is complex, centralizers are complex Lie subgroups, \mathfrak{c} is a complex vectorsubspace, and roots λ are \mathbb{C} -linear maps. None of them is pure imaginary (i.e. takes pure imaginary values on \mathfrak{c}). Thus P is empty so \mathfrak{c} is balanced. \blacksquare

5. COMPLEXIFIED CENTERS OF CENTRALIZERS IN $SL(n, \mathbb{C})$

The first step is to list the possible complexified centers $\mathfrak{c} \otimes \mathbb{C}$ and describe the root structure. This depends only on the complexified Lie algebra.

Lemma 5.1. *Let \mathfrak{g} be a real Lie algebra. Let $H \subset \mathfrak{g}$ be a subset. Let $ZZ_{\mathfrak{g}}(H)$ denote the center of its centralizer. Then*

$$ZZ_{\mathfrak{g}}(H) \otimes \mathbb{C} = ZZ_{\mathfrak{g} \otimes \mathbb{C}}(H).$$

Our approach consists in using the standard complex representation V of $\mathfrak{g} \otimes \mathbb{C}$. The root space decomposition of V under $\mathfrak{c} \otimes \mathbb{C} = ZZ_{\mathfrak{g} \otimes \mathbb{C}}(H)$ is obtained from the isotypical decomposition of V under H . This easily provides us with the decomposition of $End(V)$ under $\mathfrak{c} \otimes \mathbb{C}$, and then of the invariant subspace $\mathfrak{g} \otimes \mathbb{C} \subset End(V)$.

In this section, we treat the model case of $SL(n, \mathbb{C})$, and in the next section, the more elaborate cases $O(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$.

5.1. H -modules.

Definition 5.2. *Let H be a group. The data of a finite dimensional complex vectorspace and a homomorphism of H onto a reductive real algebraic subgroup of $Gl(V)$ is called an H -module.*

Note that H -invariant subspaces of H -modules are again H -modules. A H -module is *irreducible* if it has no proper H -submodules.

Definition 5.3. *Let $E(H)$ denote the set of equivalence classes of H -modules. Given an H -module V and $\pi \in E(H)$, let I_π , the π -isotypical component of V , denote the sum of all submodules of V belonging to the equivalence class π .*

The following Lemma is well known, but we give a full proof since it serves as a model for orthogonal, symplectic and unitary versions of it given in the next section.

Lemma 5.4. *Any H -module V splits as a direct sum of its isotypical components*

$$V = \bigoplus_{\pi \in E(H)} I_\pi.$$

Proof: Let $W \subset I_\pi$ be an irreducible invariant subspace. Since H is reductive, for each invariant subspace Z of I_π belonging to the class π , there exists an H -invariant complement to Z , thus an H -invariant projector $p_Z : I_\pi \rightarrow Z$. Since such submodules Z generate I_π , for at least one such Z , $p_Z(W) \neq 0$, thus $p_Z(W) = Z$ and W belongs to π .

More generally, if $F \subset E(H)$ is a subset, and $W \subset \sum_{\pi \in F} I_\pi$ is an irreducible invariant subspace, then W belongs to one of the classes in F . Indeed, otherwise $p_Z(W) = 0$ for all invariant subspaces Z whose class belongs to F , and these generate $\sum_{\pi \in F} I_\pi$. In particular, for every $\pi \in E(H)$, $I_\pi \cap \sum_{\pi' \neq \pi} I_{\pi'} = \{0\}$, which shows that the sum $\sum_{\pi \in E(H)} I_\pi$ is direct.

Since H is reductive, the invariant subspace $\bigoplus_{\pi \in E(H)} I_\pi$ admits an invariant complement, which contains an irreducible subspace. This is a contradiction unless $\bigoplus_{\pi \in E(H)} I_\pi = V$. \blacksquare

5.2. Centers of centralizers in $SL(n, \mathbb{C})$.

Lemma 5.5. *Let $H \subset SL(n, \mathbb{C})$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{sl}(n, \mathbb{C})$ be the center of its centralizer. Let L denote the set of nonzero roots of \mathfrak{c} in the standard representation of $\mathfrak{sl}(n, \mathbb{C})$, and d_ℓ the dimensions of the corresponding eigenspaces. Then L has $\dim(\mathfrak{c}) + 1$ elements, which satisfy exactly one linear relation,*

$$\sum_{\ell \in L} d_\ell \ell = 0.$$

Furthermore, the map

$$L \times L \setminus \text{diagonal} \rightarrow \Lambda, \quad (\ell, \ell') \mapsto \ell - \ell'$$

is one to one onto the set Λ of nonzero roots of \mathfrak{c} in the adjoint representation of $\mathfrak{sl}(n, \mathbb{C})$.

Proof: Since H is reductive, the standard action of H on \mathbb{C}^n splits into irreducibles. Let us group them into isotypical components I_ℓ : I_ℓ is the direct sum of k_ℓ isomorphic irreducible summands. According to Schur's Lemma, the group of H -automorphisms of the representation I_ℓ is isomorphic to $GL(k_\ell, \mathbb{C})$. Then

$$\begin{aligned} Z_{GL(n, \mathbb{C})}(H) &= \prod_{\ell \in L} Z_{GL(I_\ell)}(H|_{I_\ell}) \\ &\simeq \prod_{\ell \in L} GL(k_\ell, \mathbb{C}), \end{aligned}$$

whose center is $(\mathbb{C}^*)^L$, acting on \mathbb{C}^n by multiplication by a different constant on each I_ℓ . Pick a basis of \mathbb{C}^n adapted to the splitting $\mathbb{C}^n = \bigoplus_{\ell} I_\ell$. Then the center \mathfrak{c} of $Z_{SL(n, \mathbb{C})}(H)$ consists of diagonal matrices $\text{diag}(a_1, \dots, a_n)$ whose entries corresponding to basis vectors from the same I_ℓ are equal, and which sum up to 0. It follows that the elements of L generate \mathfrak{c}^* and satisfy only one linear relation, $\sum_{\ell \in L} d_\ell \ell = 0$. In particular, if (ℓ, ℓ') , (m, m')

are distinct ordered pairs of distinct elements of L , $\ell - \ell' - m + m'$ does not vanish identically on \mathfrak{c} . This shows that the map

$$L \times L \setminus \text{diagonal} \rightarrow \Lambda, \quad (\ell, \ell') \mapsto \ell - \ell',$$

which is clearly surjective onto the set of nonzero roots of the adjoint action on $\mathfrak{sl}(n, \mathbb{C})$, is injective as well. Furthermore, the root space for $\ell - \ell'$ is $\text{Hom}(I_\ell, I_{\ell'})$. \blacksquare

6. CENTRALIZERS IN ORTHOGONAL, SYMPLECTIC OR UNITARY GROUPS

In this section, the rootspace decomposition of \mathfrak{g} under the center of the centralizer of a reductive subgroup is given, when $\mathfrak{g} = \mathfrak{so}(n, \mathbb{C})$ or $\mathfrak{sp}(n, \mathbb{C})$. This is a first step in handling real forms of these Lie algebras. With little extra effort, one can treat simultaneously the case of $\mathfrak{g} = \mathfrak{su}(p, q)$. This will help treating this particular real form of $\mathfrak{sl}(n, \mathbb{C})$.

Let (V, B) be a complex vectorspace equipped with a nondegenerate binary form B of one of the following three types,

- (1) symmetric bilinear,
- (2) skew-symmetric bilinear,
- (3) symmetric sesquilinear,

which we denote by (ι, ϵ) -symmetric, $\epsilon = \pm 1$, $\iota = 1$ or $\bar{}$ (identity or conjugation). Note that the combination $(\iota, \epsilon) = (\bar{}, -1)$ makes perfect sense but does not bring anything new, since if B is a skew-symmetric sesquilinear form, then iB is symmetric sesquilinear. Let $O = O^{\iota, \epsilon}(V, B)$ denote its automorphism group (note that if B is skew-symmetric bilinear (resp. sesquilinear), this is a symplectic (resp. unitary) rather than an orthogonal group, whence the notation $O^{\iota, \epsilon}$). Let $H \subset O$ be a reductive real algebraic subgroup. In this section, we describe the center of the centralizer of H in O .

6.1. Bilinear and sesquilinear forms.

Notation 6.1. *Let ι be a continuous automorphism of \mathbb{C} , i.e. either conjugation or identity. Let $\epsilon \in \{-1, 1\}$. Let V be a complex vectorspace. A (ι, ϵ) -symmetric form on V is a real bilinear form $B : V \times V \rightarrow \mathbb{C}$ such that*

- for $\lambda, \lambda' \in \mathbb{C}$, $v, v' \in V$, $B(\lambda v, \lambda' v') = \iota(\lambda)\lambda' B(v, v')$;
- $B(v', v) = \epsilon B(v, v')$.

In other words, if $\iota = 1$ and $\epsilon = 1$, B is symmetric bilinear. If $\iota = 1$ and $\epsilon = -1$, B is skew-symmetric bilinear. If $\iota = \bar{}$ and $\epsilon = 1$, B is symmetric sesquilinear. If $\iota = \bar{}$ and $\epsilon = -1$, B is skew-symmetric sesquilinear. We shall ignore the fourth case, since if B is a skew-symmetric sesquilinear form, then iB is symmetric sesquilinear.

Notation 6.2. *1. Let ι be a continuous automorphism of \mathbb{C} , i.e. either conjugation or identity. Let V be a complex vectorspace. Then V^ι means V if $\iota = 1$, \bar{V} if $\iota = \bar{}$. Also, $V^{\iota, *}$ means the dual vectorspace V^* if $\iota = 1$, \bar{V}^* (i.e. the space of anti- \mathbb{C} -linear forms on V) if $\iota = \bar{}$.*

2. Let B be a (ι, ϵ) -symmetric form on V . Let $\sharp_B : V \rightarrow V^{\iota,*}$ denote the \mathbb{C} -linear map which maps $v \in V$ to the anti- \mathbb{C} -linear functional

$$\sharp_B(v) : v' \mapsto B(v', v).$$

Say that B is nondegenerate if \sharp_B is an isomorphism. If not, its kernel is called the nullspace of B .

Notation 6.3. Given a linear map $L : V \rightarrow V^{\iota,*}$, there is an adjoint map $L^{\iota,\top} : V \rightarrow V^{\iota,*}$ defined by

$$\langle L^{\iota,\top}(v'), v \rangle = \langle L(v), v' \rangle,$$

where $\langle v^*, v \rangle$ denotes the evaluation of a linear or anti-linear form v^* on a vector $v \in V$.

If L is ϵ -symmetric, i.e. if $L^{\iota,\top} = \epsilon L$, the formula

$$B(v, v') = \langle L(v'), v \rangle$$

defines a (ι, ϵ) -symmetric form such that $\sharp_B = L$. Therefore B and \sharp_B are equivalent data.

Example 6.4. Let W be a complex vectorspace. Then the tautological isomorphism of $V = W \times W^{\iota,*}$ to $V^{\iota,*}$ gives rise to a tautological (ι, ϵ) -symmetric form on V ,

$$(v, v^*) \cdot (w, w^*) = \epsilon \iota(\langle v^*, w \rangle) + \langle w^*, v \rangle.$$

Every two nondegenerate $(1, \epsilon)$ -symmetric forms are isomorphic. On the other hand, $(\iota, 1)$ -symmetric forms, i.e. symmetric sesquilinear forms, take real values on the diagonal, so sign and signature issues arise: two nondegenerate $(\iota, 1)$ -symmetric forms are isomorphic if and only if they have the same signature. For instance, the tautological form of Example 6.4 has vanishing signature.

6.2. Bilinear and sesquilinear H -modules.

Definition 6.5. Let H be a group. The data of a finite dimensional complex vectorspace equipped with a bilinear (either symmetric or skew-symmetric) or sesquilinear form and a homomorphism of H onto a reductive real algebraic subgroup of its automorphism group will be called a (ι, ϵ) -linear H -module.

Lemma 6.6. Let W be an irreducible H -module. The space of H -invariant bilinear (resp. sesquilinear) forms on W has dimension at most 1. A non-zero H -invariant bilinear (resp. sesquilinear) form is automatically non-degenerate, and in the bilinear case, it is either symmetric or skew-symmetric.

Proof: Let b be an H -invariant bilinear form on W . Its nullspace is H -invariant. Therefore b is either zero or non-degenerate. Assume b is nonzero and denote by $\sharp_b : W \rightarrow W^{\iota,*}$ the corresponding isomorphism (in the sesquilinear case, $\iota = \text{conjugation}$, $W^{\iota,*} = \overline{W^*}$). Let b' be an other

H -invariant bilinear form on W . Then $L = (\sharp_b)^{-1} \circ \sharp_{b'}$ is an H -equivariant endomorphism of W , thus L is a multiple of identity (Schur's Lemma). This shows that b' is a multiple of b . b has a symmetric and a skew-symmetric component. They have to be linearly dependant, in the bilinear case ($\iota = 1$), this implies that one of them vanishes. Therefore b is either symmetric or skew-symmetric. ■

Definition 6.7. *Say an H -module is (ι, ϵ) -orthogonal if it admits an invariant non-degenerate (ι, ϵ) -symmetric form.*

Corollary 6.8. *The classification of irreducible (ι, ϵ) -linear H -modules can be deduced from the classification of irreducible H -modules : the forgetful map $E^{(\iota, \epsilon)}(H) \rightarrow E(H)$ is onto, the fiber of an equivalence class of irreducible H -modules contains 1 or 2 elements depending whether it is (ι, ϵ) -orthogonal or not.*

Lemma 6.9. *Let (V, B) be a (ι, ϵ) -linear H -module. Let W and W' be distinct irreducible H -invariant subspaces. Assume that W' is not orthogonal to W . Then W' is isomorphic, as an H -module, to the (conjugate-)dual $W^{\iota, *}$ of W (in the sesquilinear case, $W^{\iota, *} = \bar{W}^*$).*

Proof: The map $v \mapsto (\sharp_B(v))|_W, W' \rightarrow W^{\iota, *}$, is H -equivariant. According to Schur's Lemma, such a map is either zero or an isomorphism, and all such maps are proportional. By assumption, it does not vanish, thus W' and $W^{\iota, *}$ are isomorphic H -modules. ■

Definition 6.10. *Let H be a group, V an H -module. Say V is bi-isotypical if there exists an irreducible H -module Z such that every irreducible invariant subspace $W \subset V$ is isomorphic either to Z or to $Z^{\iota, *}$.*

Corollary 6.11. *Let (V, B) be a non-degenerate (ι, ϵ) -linear H -module. Then V canonically splits as an orthogonal direct sum of its bi-isotypical components,*

$$V = \bigoplus_{\pi} I_{\pi, \pi^{\iota, *}},$$

where, given an equivalence class π of irreducible H -modules, $I_{\pi, \pi^{\iota, *}} = I_{\pi} + I_{\pi^{\iota, *}}$ is the sum of all irreducible invariant subspaces of V isomorphic either to π or to $\pi^{\iota, *}$. Furthermore, $I_{\pi, \pi^{\iota, *}}$ is non-degenerate, and the centralizer $Z_{O(V, B)}(H)$ of H in the automorphism group is isomorphic to a direct product,

$$Z_{O(V, B)}(H) = \prod_{\pi} Z_{O(I_{\pi, \pi^{\iota, *}})}(H).$$

6.3. Examples of bi-isotypical bilinear/sesquilinear H -modules. From now on, we analyze non-degenerate bi-isotypical (ι, ϵ) -linear H -modules. Here are two examples.

Example 6.12. Let $Z = \pi^{\oplus r}$ be an isotypical H -module. Set $V = Z \times Z^{\iota,*}$, equip it with the canonical (ι, ϵ) -symmetric form

$$(v, v^*) \cdot (w, w^*) = \epsilon \iota(\langle v^*, w \rangle) + \langle w^*, v \rangle.$$

Then the centralizer of H in $O^{(\iota, \epsilon)}(V, \cdot)$ is isomorphic to an orthogonal, symplectic or unitary group,

$$\begin{aligned} Z_{O^{(\iota, \epsilon)}(V, \cdot)}(H) &\cong O^{(\iota, \epsilon)}(2r, \mathbb{C}) \\ &:= \begin{cases} O(2r, \mathbb{C}) & \text{if } \iota = 1, \epsilon = 1, \\ Sp(2r, \mathbb{C}) & \text{if } \iota = 1, \epsilon = -1, \\ U(r, r) & \text{if } \iota = \bar{}, \epsilon = 1, \end{cases} \end{aligned}$$

if π and $\pi^{\iota,*}$ are equivalent, to a general linear group,

$$Z_{O^{(\iota, \epsilon)}(V, \cdot)}(H) \cong Gl(r, \mathbb{C})$$

otherwise.

Proof: Write $I_\pi = \bigoplus_{i=1}^r W_i$ where each W_i is H -invariant and irreducible. Fix once and for all H -isomorphisms $L_i : W_1 \rightarrow W_i$. Pick a basis of W_1 , include its images under L_i 's to get a basis of I_π . Take the image of this basis under \sharp_B to get a basis of $I_{\pi^{\iota,*}}$, this gives a basis of V in which the matrix of the bilinear/sesquilinear form B equals $\begin{pmatrix} 0 & 1 \\ \epsilon & 0 \end{pmatrix}$ (blocks have size rd where $d = \dim \pi$). In this basis, the matrix of an element g of H is diagonal in blocks of size d , with the first r diagonal blocks equal to $\pi(g)$ and the last r equal to $\pi(g^{-1})^{\iota, \top}$.

If π and $\pi^{\iota,*}$ are equivalent, endomorphisms of V which commute with H have matrices whose blocks of size d are scalar, i.e. proportional to the unit $d \times d$ matrix. In other words, they can be written $A \otimes 1$ where $A \in Gl(2r, \mathbb{C})$. Such a matrix preserves B if and only if

$$(A \otimes 1)^{\iota, \top} \left(\begin{pmatrix} 0 & 1 \\ \epsilon & 0 \end{pmatrix} \otimes 1 \right) (A \otimes 1) = \begin{pmatrix} 0 & 1 \\ \epsilon & 0 \end{pmatrix} \otimes 1,$$

i.e. iff A belongs to $O(2r, \mathbb{C})$ if $\epsilon = 1$, to $Sp(2r, \mathbb{C})$ if $\epsilon = -1$ and to $U(r, r)$ if $\iota = \text{conjugation}$. In other words,

$$Z_{O^{(\iota, \epsilon)}(V, \cdot)}(H) \cong O^{(\iota, \epsilon)}(2r, \mathbb{C}).$$

If π and $\pi^{\iota,*}$ are not equivalent, endomorphisms of V which commute with H preserve the splitting $V = I_\pi \oplus I_{\pi^{\iota,*}}$ and have matrices whose blocks of size d are scalar, i.e. proportional to the unit $d \times d$ matrix. In other words, they can be written $(A \oplus A') \otimes 1$ where $A, A' \in Gl(r, \mathbb{C})$. Such a matrix preserves B if and only if $A' = (A^{\iota, \top})^{-1}$. In other words,

$$Z_{O^{(\iota, \epsilon)}(V, \cdot)}(H) \cong Gl(r, \mathbb{C}).$$

■

Example 6.13. Let $V = \pi^{\oplus r}$ be an isotypical H -module such that π is (ι, ϵ) -orthogonal, i.e. preserves a non-degenerate (ι, ϵ) -symmetric form b . Let D be a real diagonal invertible $r \times r$ matrix. Set $B = b \otimes D$, i.e. (V, B) is an orthogonal direct sum of r real multiples of the same non-degenerate (ι, ϵ) -linear H -module. Then the centralizer of H in $O^\epsilon(V, b \otimes D)$ is isomorphic to an orthogonal/unitary group,

$$Z_{O^{\iota, \epsilon}(V, b \otimes D)}(H) \cong O^\iota(\mathbb{C}^r, D).$$

Note that in the bilinear case, $O(\mathbb{C}^r, D) = O(r, \mathbb{C})$ is a genuine orthogonal group, even when $\epsilon = -1$, i.e. when we deal with skew-symmetric forms. In the sesquilinear case, $O^\iota(\mathbb{C}^r, D) = U(p, q)$ where $p - q = \text{sign}(D)$.

Proof: Repeat the same basis of π to get a basis of V . The matrix of $B = b \otimes D$ is diagonal in blocks of size $d = \dim \pi$, with diagonal blocks equal to real multiples of b . Element $g \in H$ acts by a diagonal matrix in $d \times d$ -blocks, with diagonal blocks equal to $\pi(g)$. The centralizer of H in $Gl(V)$ consists of matrices with scalar $d \times d$ -blocks, i.e. of the form $A \otimes 1$ for $A \in Gl(r, \mathbb{C})$. (ι, ϵ) -orthogonal matrices satisfy

$$(A \otimes 1)^{\iota, \top} (b \otimes D) (A \otimes 1) = b \otimes (A^{\iota, \top} D A) = b \otimes D,$$

i.e. $A^{\iota, \top} D A = D$. Thus

$$Z_{O^\epsilon(V, B)}(H) \cong O^\iota(\mathbb{C}^r, D).$$

■

6.4. Classification of bi-isotypical bi/sesquilinear H -modules. There are 3 cases, depending whether π is (ι, ϵ) -orthogonal, $(\iota, -\epsilon)$ -orthogonal, or neither. In each case, we will need the following lemma.

Lemma 6.14. Let $V = W \oplus W'$ be a (ι, ϵ) -linear H -module where W, W' are isotropic, isotypic, and the map $L : W' \rightarrow W^{\iota, *}$, $L(v') = (\sharp_B(v'))|_W$ is an isomorphism. Then V is isomorphic to Example 6.12.

Proof: The H -map $(w + w') \mapsto (w, L(w'))$ is an isometry $W \oplus W' \rightarrow W \times W^{\iota, *}$, since

$$\begin{aligned} (v, L(v')) \cdot (w, L(w')) &= \epsilon \iota(\langle L(v'), w \rangle) + \langle L(w'), v \rangle \\ &= \epsilon \iota(\langle \sharp_B(v'), w \rangle) + \langle \sharp_B(w'), v \rangle \\ &= \epsilon \iota(B(w, v')) + B(v, w') \\ &= B(v', w) + B(v, w') \\ &= B(v + v', w + w'). \end{aligned}$$

■

Proposition 6.15. Let (V, B) be a non-degenerate bi-isotypical bilinear H -module. Assume π and $\pi^{\iota, *}$ are not isomorphic. Then (V, B) is isomorphic to Example 6.12.

Proof: Irreducible invariant subspaces of V belong either to π or $\pi^{\iota,*}$, which do not admit non-degenerate invariant bilinear/sesquilinear forms, thus all are isotropic. Lemma 6.9 implies that every two distinct irreducible invariant subspaces of I_π are orthogonal. Thus I_π is isotropic, and so is $I_{\pi^{\iota,*}}$. Since V is non-degenerate, $\sharp_B : V \rightarrow V^{\iota,*}$ induces isomorphisms $L : I_\pi \rightarrow (I_{\pi^{\iota,*}})^{\iota,*}$ and $L' : I_{\pi^{\iota,*}} \rightarrow (I_\pi)^{\iota,*}$ related by $L^\perp = \epsilon L'$. According to Lemma 6.14, this shows that V is isomorphic, as a (ι, ϵ) -linear H -module, to Example 6.12. ■

Proposition 6.16. *Let $V = I_\pi$ be an isotypical H -module equipped with a nondegenerate ϵ -symmetric bilinear form B . Assume π is $(-\epsilon)$ -orthogonal. Then (V, B) is isomorphic to Example 6.12.*

Proof: Since π is not ϵ -orthogonal, all irreducible H -submodules of V are isotropic. Let W be one of them. A H -invariant complement to W^\perp contains an irreducible H -submodule W' , which is not orthogonal to W . The H -map $L : W' \rightarrow W^*$, $v' \mapsto (\sharp_B(v'))|_W$ is non-zero, thus an isomorphism, and Lemma 6.14 implies that $W \oplus W'$ is isomorphic to Example 6.12, in particular, it is non-degenerate. Its orthogonal is again non-degenerate, isotypic, modelled on a $(-\epsilon)$ -orthogonal H -module. By induction on dimension, V is an orthogonal direct sum of copies of Example 6.12, thus isomorphic to Example 6.12. ■

Proposition 6.17. *Let $V = I_\pi$ be an isotypical H -module equipped with a nondegenerate (ι, ϵ) -symmetric bilinear form B . Assume π is (ι, ϵ) -orthogonal. Then (V, B) is isomorphic to Example 6.13.*

Proof: Let us show that V contains at least one non-degenerate irreducible invariant subspace. Pick an irreducible H -submodule W of V . If it is non-degenerate, we are done. Otherwise, W is isotropic. An H -invariant complement to W^\perp contains an irreducible H -submodule W' , which is not orthogonal to W . If W' is non-degenerate, we are done. Otherwise, W' is isotropic too. Then $W \oplus W'$ is isomorphic to $W \times W^{\iota,*}$ equipped with the canonical (ι, ϵ) -symmetric form. Indeed, the H -map $L : W' \rightarrow W^{\iota,*}$, $v' \mapsto (\sharp_B(v'))|_W$ is non-zero, thus an isomorphism, and Lemma 6.14 applies. By assumption, there exists an ϵ -symmetric H -isomorphism $M : W \rightarrow W^{\iota,*}$. Then the graph $Z = \{(w, M(w)) \mid w \in W\}$ of M is non-degenerate. Indeed, for $v, w \in W$,

$$\begin{aligned} (v, M(v)) \cdot (w, M(w)) &= \epsilon \iota(\langle M(v), w \rangle) + \langle M(w), v \rangle \\ &= 2\epsilon \Re e(\langle M(v), w \rangle) \end{aligned}$$

cannot vanish for all w , unless $v = 0$. Z is a non-degenerate irreducible (ι, ϵ) -linear H -submodule of $W \times W^{\iota,*}$, which embeds isometrically into V , this is the required subspace.

The proof of the proposition is concluded by induction on dimension. If V is irreducible, we are done. Otherwise, we just showed that V has at least one non-degenerate irreducible submodule, say W . Then the induction hypothesis applies to its orthogonal W^\perp . ■

Corollary 6.18. *Let (V, B) be a nondegenerate bi-isotypical (ι, ϵ) -linear H -module. Then (V, B) is isomorphic either to Example 6.12 or to Example 6.13. The centralizer of H in the automorphism group of (V, B) is isomorphic to a general linear group in the former case, an orthogonal group in the latter.*

6.5. Centers of centralizers in unitary groups. In this case, we directly get information on a real form.

Proposition 6.19. *Let (V, B) be a non-degenerate symmetric sesquilinear H -module. Let \mathfrak{c} denote the center of the centralizer of H in $U(V, B)$. Then root spaces of V under \mathfrak{c} correspond to isotypical components I_π under H . They fall into bi-isotypical components BI_π which are pairwise orthogonal. If $I_\pi = BI_\pi$, then the corresponding root of \mathfrak{c} is pure imaginary. If $I_\pi \neq BI_\pi$, then $BI_\pi = I_\pi \oplus I_{\bar{\pi}^*}$, the corresponding roots ℓ_π and $\ell_{\bar{\pi}^*}$ are opposite. The linear relations between roots are generated by $\ell_\pi + \ell_{\bar{\pi}^*} = 0$, $\pi \neq \bar{\pi}^*$.*

Let N be a set which contains exactly one element of each pair $\{\pi, \bar{\pi}^\}$ of equivalence classes of irreducible sesquilinear H -modules occurring in V , such that $\pi \neq \bar{\pi}^*$. Then the map $(\ell_\pi)_{\pi \in L_N} : \mathfrak{c} \rightarrow \mathbb{C}^{L_N}$ is onto.*

Proof: If $I_\pi = BI_\pi$, then, as a sesquilinear H -module, BI_π is isomorphic to Example 6.13 and contributes a unitary factor to the centralizer of H , whose center is a pure imaginary subgroup of $SL(n, \mathbb{C})$. It acts on BI_π by multiplication by a pure imaginary number. Thus BI_π is a root space for a pure imaginary root. Otherwise, BI_π is isomorphic to Example 6.12 and contributes a general linear group factor to the centralizer. Its center is a complex subgroup of $SL(n, \mathbb{C})$, it acts on I_π (resp. $I_{\bar{\pi}^*}$) by multiplication by an unrestricted complex number (resp. the opposite number). This produces a subspace \mathfrak{c}' of \mathfrak{c} which admits a complex structure, the corresponding roots are \mathbb{C} -linear and half of them (to avoid the relations $\ell_\pi + \ell_{\bar{\pi}^*} = 0$) provide complex coordinates on \mathfrak{c}' . ■

Corollary 6.20. *Let \tilde{L} be the set of roots, L_I the subset of roots which take only pure imaginary values. Then for every $\ell \in L \setminus L_I$, $-\ell$ is again a root. Let L_N be a set which contains exactly one element of each pair $\{\ell, -\ell\}$, $\ell \in L \setminus L_I$. Then*

- I_ℓ and $I_{\ell'}$ are orthogonal unless $\ell' = -\ell$;
- if $\ell \in L_I$, I_ℓ is nondegenerate;
- the map $(\ell)_{\ell \in L_N} : \mathfrak{c} \rightarrow \mathbb{C}^{L_N}$ is onto.

6.6. Centers of centralizers in ϵ -orthogonal groups. We continue our convention that

$$O^\epsilon(n, \mathbb{C}) = \begin{cases} O(n, \mathbb{C}) & \text{if } \epsilon = 1, \\ Sp(n/2, \mathbb{C}) & \text{if } \epsilon = -1. \end{cases}$$

Note that in the latter case, n has to be even.

Proposition 6.21. *Let (V, B) be a non-degenerate bilinear H -module. Let \mathfrak{c} denote the center of the centralizer of H in $O^\epsilon(V, B)$. Under \mathfrak{c} , V splits into root spaces as follows. Each isotypical component I_π where π is an irreducible H -module which is not equivalent to its contragredient π^* is a root space for a non-zero root ℓ_π . It is isotropic. The sum of all other isotypical components constitutes the 0 root space. Bi-isotypical components $I_\pi + I_{\pi^*}$ are pairwise orthogonal. Relations among non-zero roots are generated by the following*

$$\ell_\pi + \ell_{\pi^*} = 0.$$

Thus the number of non zero roots is $2\dim \mathfrak{c}$. Let L be a set containing exactly one element of each pair (π, π^) . Then $(\ell_\pi)_{\pi \in L} : \mathfrak{c} \rightarrow \mathbb{C}^L$ is a linear bijection.*

Proof: Let L' be the set of equivalence classes of irreducible H -modules which are isomorphic to their contragredient. Then

$$Z_{O^\epsilon(V, B)}(H) = \prod_{\pi \in L} GL(r_\pi, \mathbb{C}) \times \prod_{\pi \in L'} O^\epsilon(I_\pi),$$

thus

$$ZZ_{O^\epsilon(V, B)}(H) = \bigoplus_{\pi \in L} \mathbb{C}(id_{I_\pi} - id_{I_{\pi^*}}),$$

i.e. $(\ell_\pi)_{\pi \in L} : \mathfrak{c} \rightarrow \mathbb{C}^L$ is a linear bijection. Furthermore, if $\pi \in L$, the bi-isotypical component $I_\pi \oplus I_{\pi^*}$ splits into two root spaces relative to roots ℓ_π and $\ell_{\pi^*} = -\ell_\pi$. ■

Corollary 6.22. *Let \tilde{L} denote the set of roots of \mathfrak{c} on \mathbb{C}^n . The roots of \mathfrak{c} in its adjoint action on $\mathfrak{so}^\epsilon(V, B)$ are exactly all differences $\ell - \ell'$, for $\ell, \ell' \in \tilde{L}$, including 2ℓ if $\epsilon = -1$ or $\dim(I_\ell) > 1$, excluding 2ℓ if I_ℓ is 1-dimensional and $\epsilon = 1$.*

In other words, if L is a set of representatives of pairs $\{-\ell, \ell\}$ of nonzero roots, one finds 0, all sums $\pm\ell \pm \ell'$ for distinct $\ell, \ell' \in L$, sometimes $\pm 2\ell$ (depending on ϵ and $\dim(I_\ell)$), and, if 0 is also a root of \mathfrak{c} on \mathbb{C}^n , all $\pm\ell$, $\ell \in L$.

Proof: The Lie algebra $\mathfrak{so}(V, B)$ is the space of B -skew-symmetric endomorphisms of V , i.e. \mathbb{C} -linear maps $f : V \rightarrow V$ satisfying, for all $v, v' \in V$,

$$B(f(v), v') + B(v, f(v')) = 0.$$

The roots of \mathfrak{c} in the adjoint representation are differences $\lambda = \ell - \ell'$ of roots of \mathfrak{c} in \mathbb{C}^n . If $\ell' \neq \pm\ell$, the root space relative to $\ell - \ell'$ is the subspace of B -skew-symmetric elements of $Hom(I_\ell, I_{\ell'}) \oplus Hom(I_{-\ell'}, I_{-\ell})$. It does never vanish. Indeed, for every $f \in Hom(I_\ell, I_{\ell'})$, there is a unique $g \in Hom(I_{-\ell'}, I_{-\ell})$ such that (f, g) is B -skew-symmetric. The formula for g is

$$g = -\sharp_B^{-1} \circ f^\top \circ \sharp_B.$$

Here is an alternative description of $\mathfrak{so}(V, B)$: mapping $f \in \text{End}(V)$ to the bilinear form $b(v, v') = B(f(v), v')$ identifies $\text{End}(V)$ with the space $V^* \otimes V^*$ of \mathbb{C} -bilinear forms on V and $\mathfrak{so}(V, B)$ with the subspace $\Lambda^\epsilon V^*$ of $(-\epsilon)$ -symmetric \mathbb{C} -bilinear forms on V . The adjoint action of $Z \in \text{End}(V)$ is

$$Zb(v, v') = B([Z, f](v), v') = -b(v, Zv') - b(Zv, v').$$

If $v \in I_\ell$, $v' \in I_{\ell'}$ and $Z \in \mathfrak{c}$, then

$$Zb(v, v') = -(\ell + \ell')(Z)b(v, v').$$

Therefore the root space relative to 2ℓ identifies with $\Lambda^\epsilon I_\ell^*$. It vanishes if and only if $\epsilon = 1$ and $\dim(I_\ell) = 1$. \blacksquare

7. REAL FORMS OF $SL(n, \mathbb{C})$

7.1. Flexibility in $SL(n, \mathbb{R})$ and $SL(n, \mathbb{H})$.

Proposition 7.1. *Let $\mathfrak{c} \subset \mathfrak{sl}(n, \mathbb{R})$ (resp. $\mathfrak{sl}(n/2, \mathbb{H})$) be the center of the centralizer of a reductive subgroup of $SL(n, \mathbb{R})$ (resp. $SL(n/2, \mathbb{H})$). Let λ be a pure imaginary root of \mathfrak{c} in its adjoint action on $\mathfrak{sl}(n, \mathbb{R})$. Then the signature of the Killing form restricted to $\mathfrak{g}_{\lambda, \mathbb{R}}$ does not vanish. It follows that \mathfrak{c} is balanced.*

Proof: Let \mathbb{C}^n denote the standard representation of $\mathfrak{sl}(n, \mathbb{C})$. As in subsection 4.2, let $\tau(v) = \bar{v}$ in the real case, and $\tau(v) = vj$ in the quaternionic case (here, $\mathbb{C}^n = \mathbb{H}^{n/2}$ is viewed as a right quaternionic vectorspace). Let, for $f \in \mathfrak{sl}(n, \mathbb{C})$, $\sigma(f) = \tau \circ f \circ \tau^{-1}$. Then $\text{Fix}(\sigma) = \mathfrak{g} = \mathfrak{sl}(n, \mathbb{R})$ (resp. $\mathfrak{sl}(n/2, \mathbb{H})$).

Under \mathfrak{c} , \mathbb{C}^n splits into root spaces $\mathbb{C}^n = \bigoplus_\ell I_\ell$, $\dim(I_\ell) = d_\ell$. Roots are either real or come in pairs $\{\ell, \bar{\ell}\}$ (Lemma 4.1). According to Lemma 5.5, every nonzero root λ of \mathfrak{c} in its adjoint action on $\mathfrak{sl}(n, \mathbb{C})$ can be uniquely written in the form $\ell - \ell'$. Such a root is pure imaginary if and only if $\ell' = \bar{\ell}$, i.e. $\lambda = \ell - \bar{\ell}$. The corresponding root space is

$$\mathfrak{g}_\lambda = \text{Hom}(I_\ell, I_{\bar{\ell}}).$$

Let $f \in \text{Hom}(I_\ell, I_{\bar{\ell}})$, $\sigma(f) = \bar{f} \in \text{Hom}(I_{\bar{\ell}}, I_\ell)$. Pick a basis of I_ℓ and take its image by τ as a basis of $I_{\bar{\ell}}$. Let M denote the matrix of f in the chosen basis of \mathbb{C}^n . Then the matrix of $\sigma(f)$ is $\sigma(M)$, and $\text{Trace}(\sigma(f) \circ f) = \text{Trace}(\bar{M}M)$. Write $M = S + A$ where S is symmetric and A is skew-symmetric. Then

$$\text{Trace}(\bar{M}M) = \text{Trace}(\bar{S}S) + \text{Trace}(\bar{A}A) = \text{Trace}(S^*S) - \text{Trace}(A^*A)$$

has signature $\dim(\{S\}) - \dim(\{A\}) = \frac{d_\ell(d_\ell+1)}{2} - \frac{d_\ell(d_\ell-1)}{2} = d_\ell$, which is nonzero. \blacksquare

7.2. Flexibility in $SU(p, q)$.

Proposition 7.2. *Let Γ be a surface group, let $\phi : \Gamma \rightarrow SU(p, q)$ a reductive homomorphism, let \mathfrak{c} be the center of the centralizer of $\phi(\Gamma)$. Assume that \mathfrak{c} is not balanced with respect to ϕ . Then, up to conjugacy, $\phi(\Gamma)$ is contained in $S(U(p, p) \times U(q - p))$, and ϕ is maximal.*

Proof: Under \mathfrak{c} , the standard representation of $\mathfrak{sl}(n, \mathbb{C})$ splits into root spaces, $\mathbb{C}^n = \bigoplus_{\ell} I_{\ell}$. The roots of \mathfrak{c} in the adjoint representation are differences $\ell - \ell'$, and all of them indeed occur. Following Corollary 6.20, split the set of roots as $\tilde{L} = L_I \cup L_N \cup -L_N$, where L_I is the subset of pure imaginary roots. If $\ell \in L_N$, 2ℓ is a root in the adjoint representation, and it is not pure imaginary. If $\ell \in L_I$ and $\ell' \in L_N$, $\ell - \ell'$ is not pure imaginary. If ℓ and $\ell' \in L_N$, $\ell - \ell'$ is not pure imaginary either, since it factors through a surjective map $\mathfrak{c} \rightarrow \mathbb{C}^{L_N}$ and a \mathbb{C} -linear form $\mathbb{C}^{L_N} \rightarrow \mathbb{C}$. So if L_N is non empty, non pure imaginary roots span \mathfrak{c}^* , and \mathfrak{c} is balanced, contradiction. So L_N is empty, all roots on \mathbb{C}^n are pure imaginary, and the corresponding root spaces I_{ℓ} are nondegenerate and pairwise orthogonal. In the sequel, we shall replace roots by there imaginary parts without expressing it in the notation.

Each root λ in the adjoint representation can be expressed in a unique way as $\lambda = \ell - \ell'$, and

$$\mathfrak{g}_{\lambda} = \text{Hom}(I_{\ell}, I_{\ell'}).$$

A calculation shows that the signature of the natural Hermitian form on $\text{Hom}(I_{\ell}, I_{\ell'})$ equals $-\text{sign}(I_{\ell})\text{sign}(I_{\ell'})$. This time, the signature is not automatically nonzero. So different arguments, based on [4], are needed.

Assume that the sesquilinear action ρ_{λ} of Γ on \mathfrak{g}_{λ} is maximal. According to Proposition 3.2, the sesquilinear space $\text{Hom}(I_{\ell}, I_{\ell'})$ must have vanishing signature. Also, the homomorphism $Z_G(\mathfrak{c}) \rightarrow U(\text{Hom}(I_{\ell}, I_{\ell'}))$ must be tight. Lemma 3.4 implies that one of I_{ℓ} and $I_{\ell'}$ has vanishing signature and the other one is definite. Say I_{ℓ} is definite, for instance. Since compact groups do not interfere with maximality, $\text{Hom}(I_{\ell}, I_{\ell'})$ behaves like a direct sum of d_{ℓ} copies of $I_{\ell'}$, so maximality of ρ_{λ} on $\mathfrak{g}_{\lambda, \mathbb{R}}$ is equivalent to maximality of the Γ action on $I_{\ell'}$, with $T_{\lambda} = T(I_{\ell'})$. If instead $I_{\ell'}$ is definite, $T_{\lambda} = -T(I_{\ell})$.

Let D (resp. E , resp. O) denote the set of roots ℓ such that I_{ℓ} is definite (resp. has vanishing signature, resp. has non vanishing signature). As in definition 2.1, let P denote the set of roots λ such that ρ_{λ} is maximal with positive Toledo invariant. Equivalently, of differences $\ell - \ell'$, $\ell \in D$, $\ell' \in E$, such that $I_{\ell'}$ is maximal with positive Toledo invariant. Let N be the complement of $\pm P$.

Let us show that if O is non empty, \mathfrak{c} is balanced with respect to ϕ . Indeed, let $\ell_0 \in O$. Then for all $\ell' \neq \ell_0$, for $\lambda = \ell_0 - \ell'$, ρ_{λ} is not maximal, thus $\ell_0 - \ell' \in N$. Since the roots ℓ span $\mathfrak{c}^* \otimes \mathbb{C}$ and satisfy the extra equation $\sum_{\ell} d_{\ell} \ell = 0$, $\text{span}_{\mathbb{C}}(\{\ell_0 - \ell' \mid \ell' \neq \ell_0\}) = \mathfrak{c}^* \otimes \mathbb{C}$. Since $\text{span}_{\mathbb{C}}(N) = \mathfrak{c}^* \otimes \mathbb{C}$, \mathfrak{c} is balanced with respect to ϕ .

From now on, we assume that O is empty. If D or E is empty, there is no room for pairs (ℓ, ℓ') for $\ell \in D$ and $\ell' \in E$, so P is empty. Therefore we assume that both D and E are non empty. Let L_D (resp. L_E) denote the span of all differences $\ell - \ell'$ for $\ell \in D$ and $\ell' \in D$ (resp. for $\ell \in E$ and $\ell' \in E$). Then $\dim(L_D) = \text{card}(D) - 1$, $\dim(L_E) = \text{card}(E) - 1$. Since $L_D \subset \text{span}_{\mathbb{C}}(D)$, $L_E \subset \text{span}_{\mathbb{C}}(E)$ and $\text{span}_{\mathbb{C}}(D) \cap \text{span}_{\mathbb{C}}(E)$ is the line

generated by $\sum_{\ell \in D} d_\ell \ell$, $L_D \cap L_E = \{0\}$, thus $\dim(L_D + L_E) = \text{card}(D) + \text{card}(E) - 2 = \dim(\mathfrak{c}) - 1$. In the quotient space $\mathfrak{c}^* \otimes \mathbb{C}/L_D + L_E$, all elements of D (resp. of E) are mapped to the same vector ℓ_D (resp. ℓ_E), and $\ell_D \neq \ell_E$. Again, if one of the $\ell - \ell'$, $\ell \in D$, $\ell' \in E$ belongs to N , $\mathfrak{c}^* \otimes \mathbb{C}/N$ vanishes, so \mathfrak{c} is balanced with respect to ϕ , contradiction.

Therefore, all $\ell - \ell'$, $\ell \in D$, $\ell' \in E$, belong to $\pm P$. If there exists two pairs (ℓ, ℓ') , $\ell \in D$, $\ell' \in E$, such that $\Im m(\ell - \ell')$ have opposite signs in $\mathfrak{c}^*/\Im m(L_D + L_E)$, then \mathfrak{c} is balanced with respect to ϕ , contradiction.

Otherwise, all $\ell - \ell'$, $\ell \in D$, $\ell' \in E$, belong to $\pm P$ and those which belong to $+P$ project to $\mathfrak{c}^*/\Im m(L_D + L_E)$ with equal signs. This implies that the direct sum representation $\bigoplus_{\ell \in E} I_\ell$ is maximal. In other words, $\phi(\Gamma) \subset U(p, p) \times U(q - p)$ is maximal. The symmetric spaces $\mathcal{D}_{p,p}$ and $\mathcal{D}_{p,q}$ corresponding to $SU(p, p)$ and $G = SU(p, q)$ have equal ranks and the embedding $\mathcal{D}_{p,p} \hookrightarrow \mathcal{D}_{p,q}$ is isometric and holomorphic. Therefore Lemma 3.8 implies that, viewed as a homomorphism $\Gamma \rightarrow G$, ϕ is maximal as well.

■

7.3. Rigidity in $SU(p, q)$. The centralizer of $SU(p, p)$ in $SU(p, q)$, $q > p$ is $U(q - p)$ with center $\mathfrak{c} = \mathfrak{u}(1)$ generated by $Z = \begin{pmatrix} -\frac{2pi}{p+q} I_{q-p} & 0 \\ 0 & \frac{(q-p)i}{p+q} I_{2p} \end{pmatrix}$.

There is only one nonzero pair of roots $\pm i$, giving rise to the root space $\text{Hom}_{\mathbb{C}}(\mathbb{C}^{q-p}, \mathbb{C}^{2p})$. The sesquilinear form s_i on $\text{Hom}_{\mathbb{C}}(\mathbb{C}^{q-p}, \mathbb{C}^{2p})$ is, in an $SU(p, p)$ -invariant manner, the direct sum of $q - p$ copies of the $U(p, p)$ -invariant Hermitian form on \mathbb{C}^{2p} . Therefore the corresponding Toledo invariant is equal to $q - p$ times the Toledo invariant obtained for $q = p + 1$. In this case, the centralizer of \mathfrak{c} is $U(1) \times SU(p, p)$, acting on the real root space via the standard complex representation of its second factor. The representation ρ_i induced on a surface subgroup $\Gamma \subset U(1) \times SU(p, p)$ is maximal if and only if the projection of Γ to $SU(p, p)$ is maximal. Thus a maximal surface subgroup of $SU(p, p)$ is not flexible in $SU(p, q)$, $q > p$, as is well known. Such subgroups exist (Theorem 1.8) and are known to be automatically discrete, [1], [3].

8. REAL FORMS OF $O(n, \mathbb{C})$ AND $Sp(n, \mathbb{C})$

8.1. Non pure imaginary roots.

Lemma 8.1. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. If one of the roots of \mathfrak{c} on \mathbb{C}^n is not pure imaginary, then \mathfrak{c} is balanced.*

Proof: By contradiction. Assume that \mathfrak{c} is not balanced and at least one of the roots of \mathfrak{c} on \mathbb{C}^n is not pure imaginary. Consider the torus $\mathfrak{c} \otimes \mathbb{C} \subset \mathfrak{so}^\epsilon(n, \mathbb{C})$ and the set $L \cup -L$ of its nonzero roots on \mathbb{C}^n (Proposition 6.21). Write $L = L_I \cup L_N$ where $L_I \subset L$ denotes the set of roots whose restriction to \mathfrak{c} takes only pure imaginary values and L_N its complement. By assumption, L_N is nonempty.

Assume first that L_I is nonempty too. None of the roots $\lambda = \pm\ell \pm \ell'$, $\ell \in L_I$, $\ell' \in L_N$, is pure imaginary, so none of them belongs to $\pm P$. They generate $\text{span}(L) = \mathfrak{c}^*$, thus \mathfrak{c} is balanced, contradiction. Therefore, $L = L_N$.

Let $v \in \mathfrak{c}$ be such that $\lambda(v) = 0$ for all roots λ of \mathfrak{c} in the adjoint representation which are not pure imaginary. For every distinct ℓ and $\ell' \in L$, one of $\ell - \ell'$ and $\ell + \ell'$ is not pure imaginary, thus $\ell'(v) = \pm\ell(v)$.

Assume that there exists $\ell \in L$ such that ℓ or 2ℓ is a root, then $\ell(v) = 0$ as well, which implies that $\ell'(v) = 0$ for all $\ell' \in L$, and $v = 0$. In other words, in that case, non pure imaginary roots span \mathfrak{c}^* , and \mathfrak{c} is balanced, contradiction. Therefore,

- 0 is not a root of \mathfrak{c} on \mathbb{C}^n (i.e., all root spaces correspond to elements of $\pm L$);
- for every root $\ell \in L$, 2ℓ is not a root of \mathfrak{c} in the adjoint representation.

This implies that $\epsilon = 1$ and all root spaces I_ℓ have dimension 1, i.e. $\mathfrak{c} \otimes \mathbb{C}$ is a maximal torus of $\mathfrak{so}(n, \mathbb{C})$. Its centralizer in $\mathfrak{so}(n, \mathbb{C})$ is abelian, so are $Z_G(\mathfrak{c})$ and H , up to finite index. A homomorphism from an abelian group cannot be tight, so no symplectic action on root spaces can be maximal, P is empty, and \mathfrak{c} is balanced again, contradiction. \blacksquare

8.2. List of real forms. Let us denote by B the matrix $\begin{pmatrix} 0 & I_m \\ \epsilon I_m & 0 \end{pmatrix}$ or $\begin{pmatrix} 0 & I_m & 0 \\ I_m & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ (depending whether n is even or odd). The real forms of $O^\epsilon(n, \mathbb{C}) = O^\epsilon(\mathbb{C}^n, B)$ are

- (1) For n even, $\epsilon = -1$, $Sp(n/2, \mathbb{R})$, consisting of matrices with real entries in $O^{-1}(n, \mathbb{C})$. It is the fixed point set of complex conjugation.
- (2) When $p \leq q$, $p + q = n$, $O(p, q)$ is the fixed point set of $M \mapsto E\bar{M}E$,

$$\text{where } E = \begin{pmatrix} I_p & 0 & 0 & 0 \\ 0 & 0 & 0 & I_{m-p} \\ 0 & 0 & I_p & 0 \\ 0 & I_{m-p} & 0 & 0 \end{pmatrix} \text{ if } n = 2m \text{ is even, and } E = \begin{pmatrix} I_p & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & I_{m-p} & 0 \\ 0 & 0 & I_p & 0 & 0 \\ 0 & I_{m-p} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \text{ if } n = 2m + 1 \text{ is odd.}$$

- (3) When $p \leq q$, $2p + 2q = n$, $\epsilon = -1$, $Sp(p, q)$ is the fixed point set of $M \mapsto e(M^{-1})^*e$ where $e = \begin{pmatrix} I_p & 0 & 0 & 0 \\ 0 & -I_q & 0 & 0 \\ 0 & 0 & -I_q & 0 \\ 0 & 0 & 0 & I_p \end{pmatrix}$.

- (4) For n even, $SO^*(n)$, sometimes denoted by $SO(n/2, \mathbb{H}) = O(n, \mathbb{C}) \cap Gl(n/2, \mathbb{H})$, is the fixed point set of $M \mapsto JM\bar{J}$ where $J = \text{diag}(j, \dots, j)$ et $j = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

We observe that in each case, there exists a complex vectorspace V , an anti- \mathbb{C} -linear map τ of V such that $\tau^{-1} = \eta\tau$, $\eta = \pm 1$, and a τ -equivariant nondegenerate ϵ -symmetric bilinear form B on V such that $G = \{f \in O^\epsilon(V, B); \sigma(f) = f\}$, where the real structure σ takes the form

$$\sigma(f) = \tau \circ f \circ \tau^{-1}.$$

τ -equivariance means that

$$B(\tau(v), \tau(v')) = \overline{B(v, v')}.$$

For the two former families, start with a real orthogonal or symplectic vectorspace (\mathbb{R}^n, b) , let $V = \mathbb{R}^n \otimes \mathbb{C}$, $B = b \otimes \mathbb{C}$ and τ is conjugation, $\eta = 1$.

For the two latter, start with a right quaternionic vectorspace $\mathbb{H}^{n/2}$, pick a basis $(1, i, j, k)$ of \mathbb{H} , view $V = \mathbb{H}^{n/2}$ with right multiplication by i as a complex vectorspace. Let τ be right multiplication by j , so that $\eta = -1$, and B be a direct sum of multiples of the following bilinear forms on \mathbb{H} ,

$$B(q, q') = \begin{cases} j\bar{q}q' - ij\bar{q}q'i & \text{if } \epsilon = -1, \\ j\bar{q}jq' - ij\bar{q}jq'i & \text{if } \epsilon = 1. \end{cases}$$

8.3. The sesquilinear structure.

Proposition 8.2. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer.*

- (1) *Let ℓ be a pure imaginary root of \mathfrak{c} on \mathbb{C}^n . Let I_ℓ denote the corresponding root space. Then I_ℓ inherits a $Z_G(\mathfrak{c})$ -invariant nondegenerate sesquilinear form s_ℓ .*
- (2) *Assume that all roots of \mathfrak{c} on \mathbb{C}^n are pure imaginary. Then the centralizer $Z_G(\mathfrak{c})$ of \mathfrak{c} in G is a product of the fixator of the orthogonal of the 0 root space I_0 and of unitary groups,*

$$Z_G(\mathfrak{c}) \cong G_{I_0^\perp} \times \prod_{\ell \in L} U(I_\ell, s_\ell).$$

Proof: 1. Let $s_\ell(v, v') = B(\tau(v), v')$. This is an $\eta\epsilon$ -symmetric sesquilinear form. It is nondegenerate on I_ℓ because B is and all root spaces but $I_{-\ell} = \tau(I_\ell)$ are B -orthogonal to I_ℓ . If $g \in Z_G(\mathfrak{c})$, i.e g belongs to G and commutes with \mathfrak{c} , then g leaves all root spaces $I_{\pm\ell}$ invariant, it commutes with τ , and is isometric for B , thus it is isometric for s_ℓ .

2. Let $g \in G$ commute with \mathfrak{c} . Let $g_0 \in GL(n, \mathbb{C})$ be the element with coincides with g on I_0 and fixes the sum of all I_ℓ , $\ell \in \pm L$, i.e. I_0^\perp . Then $g_0 \in G$. Indeed, g_0 commutes with τ and preserves B . Also, restrict g to each I_ℓ , $\ell \in L$. This yields an injective homomorphism $Z_G(\mathfrak{c}) \rightarrow G_{I_0^\perp} \times \prod_{\ell \in L} U(I_\ell, s_\ell)$.

Conversely, let $g_0 \in G$ be an element which fixes all I_ℓ . For each $\ell \in L$, pick an element $g_\ell \in U(I_\ell, s_\ell)$, extend it to $I_{-\ell}$ so that $g_\ell = \tau \circ g_\ell \circ \tau^{-1}$, extend this map trivially to other $I_{\ell'}$ and to I_0 . The obtained linear map belongs to $Z_G(\mathfrak{c})$. Indeed, it preserves B , commutes with τ and preserves each I_ℓ . Multiplying g_0 with the g_ℓ 's yields the inverse isomorphism $G_{I_0^\perp} \times \prod_{\ell \in L} U(I_\ell, s_\ell) \rightarrow Z_G(\mathfrak{c})$. \blacksquare

Lemma 8.3. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Let ℓ, ℓ' be distinct pure imaginary roots of \mathfrak{c} on \mathbb{C}^n , and $\lambda = \ell - \ell'$ the corresponding pure imaginary root of \mathfrak{c} in its adjoint action on \mathfrak{g} . Let $I_\ell, I_{\ell'}$ and \mathfrak{g}_λ denote the corresponding root spaces.*

1. *If $\ell' \neq -\ell$, then, as a sesquilinear $Z_G(\mathfrak{c})$ -module, \mathfrak{g}_λ is isomorphic to $\text{Hom}(I_\ell, I_{\ell'})$ equipped with the natural sesquilinear form*

$$(f, f') \mapsto \text{Trace}(f^* \circ f'),$$

where $f^* \in \text{Hom}(I_{\ell'}, I_\ell)$ is the adjoint of f with respect to the sesquilinear forms s_ℓ and $s_{\ell'}$.

2. *If $\ell' = -\ell$, then, as a sesquilinear $Z_G(\mathfrak{c})$ -module, \mathfrak{g}_λ is isomorphic to the subspace of skew-symmetric forms in the space $I_\ell^* \otimes I_\ell^*$ of \mathbb{C} -bilinear forms on I_ℓ , equipped with its natural sesquilinear form*

$$(b, b') \mapsto \text{Trace}((\sharp_{s_\ell})^{-1} \circ (\overline{\sharp_b})^\top \circ (\sharp_{s_\ell})^{-1} \circ \sharp_{b'}).$$

Proof: 1. Assume first that ℓ and ℓ' are linearly independent. According to Corollary 6.22, \mathfrak{g}_λ is the space of B -skew-symmetric elements of $\text{Hom}(I_\ell, I_{\ell'}) \oplus \text{Hom}(I_{-\ell'}, I_{-\ell})$. Given $f \in \text{Hom}(I_\ell, I_{\ell'})$ and $g \in \text{Hom}(I_{-\ell'}, I_{-\ell})$, $X = (f, g)$ is B -skew-symmetric means that for all $v \in I_\ell$ and $w \in I_{-\ell'}$,

$$B(f(v), w) + B(v, g(w)) = 0.$$

Since \sharp_B identifies $I_{-\ell}$ with the dual of I_ℓ , given $f \in \text{Hom}(I_\ell, I_{\ell'})$, there exists a unique $g \in \text{Hom}(I_{-\ell'}, I_{-\ell})$ such that $X = (f, g)$ is B -skew-symmetric. This shows that, as a $Z_G(\mathfrak{c})$ -module, $\mathfrak{g}_{\ell-\ell'}$ is isomorphic to $\text{Hom}(I_\ell, I_{\ell'})$.

Since $\sigma(X) = (\tau \circ g \circ \tau^{-1}, \tau \circ f \circ \tau^{-1})$,

$$\sigma(X) \circ X = \eta(\tau \circ g \circ \tau \circ f, \tau \circ f \circ \tau \circ g),$$

therefore

$$\begin{aligned} \text{Trace}(\sigma(X) \circ X) &= \eta \text{Trace}(\tau \circ g \circ \tau \circ f) + \eta \text{Trace}(\tau \circ f \circ \tau \circ g) \\ &= 2\eta \text{Trace}(\tau \circ g \circ \tau \circ f). \end{aligned}$$

If $v \in I_\ell$,

$$\begin{aligned}
s_\ell(v, \tau \circ g \circ \tau \circ f(v)) &= B(\tau(v), \tau \circ g \circ \tau \circ f(v)) \\
&= \overline{B(v, g \circ \tau \circ f(v))} \\
&= \overline{-B(f(v), \tau \circ f(v))} \\
&= -\epsilon \overline{B(\tau \circ f(v), f(v))} \\
&= -\epsilon \overline{s_\ell(f(v), f(v))} \\
&= -\epsilon \overline{s_\ell(v, f^* \circ f(v))}.
\end{aligned}$$

Summing over an orthogonal basis for s_ℓ yields

$$\text{Trace}(\tau \circ g \circ \tau \circ f) = -\epsilon \text{Trace}(f^* \circ f),$$

hence the Killing sesquilinear form on \mathfrak{g}_λ is proportional to $\text{Trace}(f^* \circ f)$.

2. Let $f \in \text{Hom}(I_\ell, I_{-\ell})$. Then f belongs to $\mathfrak{so}(V, B)$ (f is B -skew-symmetric) if and only if $f^* = -\sigma(f)$. Indeed,

$$\begin{aligned}
&\forall v, w \in I_\ell, \quad B(f(v), w) + B(v, f(w)) = 0 \\
&\Leftrightarrow \forall v, w \in I_\ell, \quad s_{-\ell}(f(v), \tau(w)) + s_\ell(v, \tau \circ f(w)) = 0 \\
&\Leftrightarrow \forall v \in I_\ell, \forall v' \in I_{-\ell}, \quad s_{-\ell}(f(v), v') + s_\ell(v, \tau \circ f \circ \tau^{-1}(v')) = 0 \\
&\Leftrightarrow f^* = -\tau \circ f \circ \tau^{-1}.
\end{aligned}$$

Thus, for B -skew-symmetric f ,

$$\text{Trace}(\sigma(f) \circ f) = \text{Trace}(\tau \circ f \circ \tau^{-1} \circ f) = -\text{Trace}(f^* \circ f).$$

Now f is B -skew-symmetric if and only if the bilinear form $b(v, v') = B(f(v), v')$ on I_ℓ is $(-\epsilon)$ -symmetric. Since $\sharp_B = f^\top \circ \sharp_B = \eta f^\top \circ \tau^\top \circ \sharp_{s_\ell}$ and $f^* = (\sharp_{s_\ell})^{-1} \circ (\overline{f})^\top \circ \sharp_{s_\ell}$,

$$f^* \circ f' = \eta (\sharp_{s_\ell})^{-1} \circ (\overline{\sharp_b})^\top \circ (\sharp_{s_\ell})^{-1} \circ \sharp_{b'}.$$

■

Lemma 8.4. *Let V be a n -dimensional complex vectorspace equipped with a nondegenerate sesquilinear form of signature s . Then the signatures of the induced sesquilinear forms on $V^* \otimes V^*$, S^2V^* and Λ^2V^* are equal to s^2 , $(s^2 + n)/2$ and $(s^2 - n)/2$ respectively.*

Proof: Fix a basis of V . If D denotes the matrix of the sesquilinear form in this basis, and b, b' the matrices of two bilinear forms on V , the induced sesquilinear form on $V^* \otimes V^*$ is

$$\text{Trace}(\overline{b}^\top D^{-1} b' D^{-1}).$$

One can assume that S is diagonal with entries $d_m = 1$ (p times) and -1 (q times), $n = p + q$, $s = p - q$. Then

$$\text{Trace}(\overline{b}^\top D^{-1} b' D^{-1}) = \sum_{m, m'} d_m d_{m'} |b_{mm'}|^2$$

is diagonal again. The signature of the whole space $V^* \otimes V^*$ is $\sum_{m, m'} d_m d_{m'} = s^2$. The signature of the subspace $\Lambda^2 V^*$ of skew-symmetric forms is

$$\sum_{m < m'} d_m d_{m'} = \frac{p(p-1)}{2} + \frac{q(q-1)}{2} - pq = \frac{s^2 - n}{2}.$$

The signature of the subspace $S^2 V^*$ of symmetric forms is

$$\sum_{m \leq m'} d_m d_{m'} = n + \sum_{m < m'} d_m d_{m'} = \frac{s^2 + n}{2}.$$

■

Lemma 8.5. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $\phi : \Gamma \rightarrow G$ be a reductive homomorphism. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Let ℓ, ℓ' be distinct nonzero pure imaginary roots of \mathfrak{c} on \mathbb{C}^n , and $\lambda = \ell - \ell'$ the corresponding pure imaginary root of \mathfrak{c} in its adjoint action on \mathfrak{g} .*

- (1) *If $\ell' \neq -\ell$, and $\lambda \in \pm P$, then one of s_ℓ and $s_{\ell'}$ has vanishing signature and the other is definite.*
- (2) *If $\ell' = -\ell$, then $\lambda = 2\ell$ either is not a root or does not belong to $\pm P$.*

Proof: If $\lambda \in P$, \mathfrak{g}_λ has vanishing signature. So does $U(\text{Hom}(I_\ell, I_{\ell'}))$ (resp. $U(\Lambda^\epsilon(I_\ell))$) with its natural sesquilinear form, according to Lemma 8.3. In particular, these groups are of tube Hermitian type. Furthermore, the $Z_G(\mathfrak{c})$ action on \mathfrak{g}_λ is tight. According to Proposition 8.2, $Z_G(\mathfrak{c})$ is a product of groups.

1. If $\ell' \neq -\ell$, among the factors, only $U(I_\ell, s_\ell)$ and $U(I_{\ell'}, s_{\ell'})$ act non trivially on \mathfrak{g}_λ , thus the morphism

$$U(I_\ell) \times U(I_{\ell'}) \rightarrow U(\text{Hom}(I_\ell, I_{\ell'}))$$

must be tight. Lemma 3.4 applies and one of the left hand groups is compact and the other has vanishing signature.

2. If $\ell' = -\ell$, among the factors, only $U(I_\ell, s_\ell)$ acts non trivially on \mathfrak{g}_λ , thus the morphism

$$U(I_\ell) \rightarrow U(\Lambda^\epsilon(I_\ell))$$

must be tight. Lemma 3.3 implies that $U(I_\ell)$ is of tube type, so s_ℓ has vanishing signature. Lemma 8.4 shows that the signature of $\Lambda^\epsilon(I_\ell)$ is $\pm \dim(I_\ell)/2$, which does not vanish, contradiction. We conclude that if 2ℓ is a root, it does not belong to $\pm P$. ■

Proposition 8.6. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $\phi : \Gamma \rightarrow G$ be a reductive homomorphism. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Then \mathfrak{c} is balanced with respect to ϕ unless $\epsilon = 1$, $\dim(\mathfrak{c}) = 1$, \mathfrak{c} is generated by a root ℓ with a 1-dimensional root space I_ℓ , the sesquilinear form on I_0 has vanishing signature, G_{I_0} is reductive Hermitian of tube type and tightly embedded in $U(I_0)$, and the homomorphism $\Gamma \rightarrow U(I_0)$ is maximal.*

Proof: Assume that \mathfrak{c} is not balanced. From Lemma 8.1, we know that roots have to be pure imaginary. We take their imaginary parts without mentioning it explicitly.

If $\epsilon = -1$ or if all I_ℓ , $\ell \in L$, have dimension > 1 , then all 2ℓ are roots and do not belong to $\pm P$, so they all belong to the complement N of P (Lemma 8.5). Since they generate \mathfrak{c}^* , \mathfrak{c} is balanced, contradiction. So $\epsilon = 1$ and the set D of roots $\ell \in L$ such that $\dim(I_\ell) = 1$ is nonempty.

Assume that D has at least 2 elements. According to Lemma 8.5, combinations $\pm\ell \pm \ell'$, for $\ell, \ell' \in D$, belong to N . Since they span $\text{span}(D)$, and multiples $2\ell''$, $\ell'' \notin D$ span $\text{span}(L \setminus D)$, N spans \mathfrak{c}^* and \mathfrak{c} is balanced, contradiction. So D has exactly one element, denoted by ℓ_0 .

Assume that $L \neq D$. Since $\dim(\mathfrak{c}^*/\text{span}(N)) = 1$, for all $\ell \in L \setminus D$, all combinations $\pm\ell_0 \pm \ell$ belong to $\pm P$. Since 0 does not belong to the convex hull of the image of P in $\mathfrak{c}^*/\text{span}(N)$, this convex hull contains exactly one of ℓ_0 and $-\ell_0$, say ℓ_0 . Then $\ell_0 - \ell$ and $\ell_0 + \ell$ belong to P . Thus the sesquilinear representation of Γ in $\text{Hom}(I_{\ell_0}, I_\ell \oplus I_{-\ell})$ is maximal. Lemma 3.5 allows to replace $\text{Hom}(I_{\ell_0}, I_\ell \oplus I_{-\ell})$ with $I_\ell \oplus I_{-\ell}$. However, as a sesquilinear vectorspace, $I_{-\ell}$ is isomorphic to $\overline{I_{-\ell}}$, so, with Lemma 3.6,

$$T(I_\ell \oplus I_{-\ell}) = T(I_\ell) + T(I_{-\ell}) = T(I_\ell) - T(I_\ell) = 0,$$

contradicting maximality. So $L = D$ and $\dim(\mathfrak{c}) = 1$.

Assume that 0 is not a root of \mathfrak{c} on \mathbb{C}^n . Then $\dim(V) = 2$ and $Z_G(\mathfrak{c}) \cong U(1)$, which cannot have any maximality property. So 0 is a root on \mathbb{C}^n , $\pm\ell$ are roots of \mathfrak{c} in the adjoint representation. Since 0 does not belong to the convex hull of P in \mathfrak{c}^* , exactly one of $\pm\ell_0$ belongs to P , say $\ell_0 \in P$. Then $\text{Hom}(I_{\ell_0}, I_0)$ is a maximal sesquilinear representation of Γ , so does I_0 , by Lemma 3.5. In particular, the signature of (I_0, s_0) vanishes, i.e. $U(I_0)$ has tube type. Also, the morphism $Z_G(\mathfrak{c}) \rightarrow PU(I_0)$ is tight. $Z_G(\mathfrak{c})$ acts on I_0 via the group $G_{I_0^\perp}$, so the injection $G_{I_0^\perp} \rightarrow PU(I_0)$ is tight. This implies that $G_{I_0^\perp}$ is reductive Hermitian (Theorem 7.1 of [4]), and that $G_{I_0^\perp}$ has tube type (Lemma 3.3). \blacksquare

Corollary 8.7. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $\phi : \Gamma \rightarrow G$ be a reductive homomorphism. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Then \mathfrak{c} is balanced with respect to ϕ unless $G = SO^*(2n)$, n odd, and $\phi(\Gamma) \subset SO^*(2n-2) \times SO^*(2)$ is maximal.*

Proof: In view of Proposition 8.6, there merely remains to determine which pairs $(G, G_{I_0^\perp})$ can lead to unbalanced centralizers, when G is a real form of $O(N, \mathbb{C})$, i.e. $G = O(P, Q)$ or $G = SO^*(2n)$.

In the real case, I_0 is real, $G_{I_0^\perp}$ is a real orthogonal group $O(p, q)$, which tightly injects into $U(I_0, s_0) = U(p, q)$. We also know that the signature $p - q$ of s_0 vanishes. $O(p, q)$ is reductive Hermitian only if $p = 2$. But Lemma 3.7 states that the inclusion $O(2, 2) \hookrightarrow U(2, 2)$ is not tight. So \mathfrak{c} is always balanced if $G = O(P, Q)$ is a real orthogonal group.

In the quaternionic case, I_0 is quaternionic and carries a nondegenerate $(-, -1)$ -binary form, $G_{I_0^\perp}$ is a quaternionic skew-unitary group $SO^*(2p)$, $p = \dim_{\mathbb{H}}(I_0)$. Since its centralizer in $G = SO^*(2n)$ is compact and 1-dimensional, $n = p + 1$. Since $G_{I_0^\perp}$ has tube type, p is even and n is odd. The homomorphism $\Gamma \rightarrow U(I_0) \cong U(n, n)$ is maximal.

.....

hence, viewed as a homomorphism $\Gamma \rightarrow G_{I_0^\perp}$, ϕ is maximal. The symmetric spaces \mathcal{G}_p and \mathcal{G}_{p+1} corresponding to $G_{I_0^\perp}$ and G have equal ranks and the embedding $\mathcal{G}_p \hookrightarrow \mathcal{G}_{p+1}$ is isometric and holomorphic. Therefore Lemma 3.8 implies that, viewed as a homomorphism $\Gamma \rightarrow G$, ϕ is maximal as well.

■

9. PROOF OF THEOREM 1

For homomorphisms $\Gamma \rightarrow G$ with reductive Zariski closure, the proof of Theorem 1 follows from Theorem 2.2, the classification of classical simple Lie groups and the case by case analysis of balancedness in sections 7 and 8.

Here is how the problem is reduced to the case of reductive homomorphisms. Theorem 2 of [12] asserts that if $\text{genus}(\Gamma) \geq 2\dim(G)^2$ and G is semisimple, the space $\text{Hom}(\Gamma, G)$ falls into two types of connected components: in some of them, Zariski dense homomorphisms are dense; others do not contain any Zariski dense homomorphism (call them rigid). Let $\phi : \Gamma \rightarrow G$ be an arbitrary homomorphism. If ϕ cannot be approximated by Zariski dense homomorphisms, then ϕ belongs to a rigid component \mathcal{C} . Proposition 8.3 of [12] asserts that \mathcal{C} contains a reductive homomorphism ψ . The reductive case of Theorem 1 implies that G is Hermitian of non tube type and ψ is maximal. Since Toledo invariants are constant on connected components of $\text{Hom}(\Gamma, G)$, ϕ is maximal as well.

REFERENCES

- [1] Steven B. Bradlow, Oscar García-Prada and Peter B. Gothen, Surface Group Representations and $U(p, q)$ -Higgs Bundles, *J. Differen. Geom.* **64**, (2003), 111–170.
- [2] Steven B. Bradlow, Oscar García-Prada and Peter B. Gothen, Maximal surface group representations in isometry groups of classical Hermitian symmetric spaces. *Geom. Dedicata* **122**, (2006), 185–213.
- [3] Marc Burger, Alessandra Iozzi, Anna Wienhard, Surface group representations with maximal Toledo invariant. arXiv:math/0605656.
- [4] Marc Burger, Alessandra Iozzi, Anna Wienhard, Tight homomorphisms and Hermitian symmetric spaces. arXiv:math/0710.5641
- [5] Jean-Louis Clerc and Bent Ørsted, The Gromov norm of the Kaehler class and the Maslov index. *Asian J. Math.* **7**, (2003), 269–295.
- [6] Antun Domic and Domingo Toledo, The Gromov norm of the Kaehler class of symmetric domains. *Math. Ann.* **276** (1987), 425–432.

- [7] William Goldman, Discontinuous groups and the Euler class. Thesis, University of California at Berkeley (1980).
- [8] William Goldman, Representations of fundamental groups of surfaces. *Geometry and topology* (J. Alexander and J. Harer, Eds), *Lect. Notes Math.* **1167**, Springer, 1985, 95–117.
- [9] William Goldman and John Millson, Local rigidity of discrete groups acting on complex hyperbolic space. *Invent. Math.* **88** (1987), 495–520.
- [10] William Goldman and John Millson, The deformation theory of representations of fundamental groups of compact Kähler manifolds. *Publ. Math. I.H.É.S.* **67** (1988), 43–96.
- [11] Luis Hernández Lamóneda, Maximal representations of surface groups in bounded symmetric domains. *Trans. Amer. Math. Soc.* **324**, (1991), 405–420.
- [12] Inkang Kim and Pierre Pansu, Density of Zariski density for surface groups. [arXiv:math/](https://arxiv.org/abs/math/)
- [13] John Milnor, On the existence of a connection with curvature zero. *Comment. Math. Helv.* **32**, (1958), 215–223.
- [14] Ichirô Satake, Algebraic structures of symmetric domains. *Kanô Memorial Lectures*, **4**. Iwanami Shoten, Tokyo; Princeton University Press, Princeton, N.J., (1980).
- [15] Domingo Toledo, Representations of surface groups in complex hyperbolic space, *J. Diff. Geom.* **29** (1989), 125–133.
- [16] Vladimir Turaev, A cocycle of the symplectic first Chern class and Maslov indices. *Funktsional. Anal. i Prilozhen.* **18**, (1984), 43–48.

1991 *Mathematics Subject Classification.*51M10, 57S25.

Key words and phrases. Algebraic group, symmetric space, rigidity, group cohomology, moduli space

Inkang Kim
 School of Mathematics
 KIAS, Heogiro 87, Dongdaemen-gu
 Seoul, 130-722, Korea
inkang@kias.re.kr

Pierre Pansu
 Département de Mathématiques et applications
 UMR 8553 du CNRS
 École Normale Supérieure
 45 rue d'Ulm
 75230 Paris Cédex 05, France
pierre.pansu@math.u-psud.fr